

# Maggot debridement therapy: Utility in chronic wounds and a perspective beyond

Sig AK, Koru O & Araz E

## ABSTRACT

Complementary medicinal techniques have a wide history, but the recent focus of modern medicine has been on possible effect mechanisms. Chronic wounds are not only a problem to their patients, but also a burden on the healthcare system due to their extensive costs.

Among complementary therapies, maggot debridement therapy (MDT) is by far the most widely studied method in the field of wound prevention and management. MDT is a very effective treatment technique for chronic wounds and, in addition, is a cheap and easy-to-use method, with only minor, rare adverse effects. Many studies have attempted to clarify the actual effect mechanisms of MDT. MDT works in four ways: debridement, antimicrobial effects, wound healing, and biofilm degradation. These actions are mainly achieved by protein substances exhibiting very variable molecular masses and actions.

MDT has a certain positive effect on chronic wounds and is a strong candidate to maintain a bold presence in

the multidisciplinary approach to chronic wound care. Its cost, application simplicity, minimal side effects, and easy-accessibility are major superiorities among other wound-care methods.

*Keywords:* *Lucilia sericata, chronic wound care, larval debridement, biosurgery.*

## INTRODUCTION

Complementary medicinal techniques used to be controversial applications for medical professionals. Recently, however, multiple studies indicating the probable benefits of these techniques to medicine have widely changed this opinion in a positive way<sup>1-3</sup>. One of these methods, maggot debridement therapy (MDT; also called larval therapy or biosurgery), is the most widely studied complementary medicinal technique in the field of wound prevention and management, and has been included in routine medical applications in many countries worldwide<sup>4</sup>. MDT has a bolder presence among these techniques, as it has been evaluated by many scientific research studies, and its medicinal effects have been observed<sup>4-7</sup>. The mechanisms of MDT action have not been totally examined, but it seems there is a combined and bound mechanistic circle influenced by the maggot itself, patient immunity, wound type, and the infective microorganisms.

Although MDT has recently been accepted worldwide, its usage has an extensive history. While some reports indicate the usage of MDT before 1900, most medical studies using MDT have been completed in the 1900s. As chronic wounds have become more relevant, cheap, effective, and easily applicable methods have been under investigation. To this effect, many studies on MDT have been published<sup>5-9</sup>.

Many larval types have been investigated for MDT, but *Lucilia sericata* larvae are the most widely studied and used maggots. The application procedure (free-range, biobag) may slightly change the strength of activity, and it is chosen depending on clinical conditions, wound aetiology, and physicians' opinion. The effects of MDT on venous stasis ulcers, pressure wounds, neuropathic ulcers (diabetic foot ulcers), traumatic, and post-surgical non-healing wounds have been previously investigated, and results were very promising<sup>4,10,11</sup>.

### Ali Korhan Sig\*

MD

Hacettepe University, Faculty of Medicine,  
Department of Medical Microbiology  
Ankara, Turkey

Email dr\_korhan@hotmail.com

### Ozgur Koru

MD

Health Sciences University, Gulhane Medical  
School, Department of Medical Microbiology,  
Ankara, Turkey

### Engin Araz

MD, PhD

Health Sciences University, Gulhane Medical  
School, Department of Medical Microbiology,  
Ankara, Turkey

\* Corresponding author

Although many researchers have particularly focused on effect mechanisms, there is still a long way to go to total understanding. The effect mechanism of MDT can be divided into four main titles: debridement, antimicrobial effects, wound healing, and biofilm degradation effect<sup>4,10</sup>. These titles are tightly related, and readers should keep in mind that these mechanisms are like a chain reaction, and cannot be evaluated separately. We, herein, review MDT focusing on application, effect mechanisms and clinical indications.

## CHRONIC WOUNDS AND WOUND DRESSINGS

Chronic wounds are practically defined as wounds that show an unhealed condition in three months' time. Many aetiologies cause chronic wounds, and they are usually classified into four groups: venous stasis ulcers, ischaemic wounds, diabetic foot ulcers, and pressure injuries (ulcers)<sup>12</sup>.

In Western countries, nearly 1% of the population has a chronic wound, and chronic wound care is a serious economic burden on the healthcare system<sup>12</sup>. In the cost modelling of Graves *et al.*<sup>13</sup>, it was reported that pressure injuries, diabetic foot ulcers, venous ulcers and artery insufficiency ulcers caused a total annual expense of approximately US\$1.65 billion, US\$250 million, US\$803 million and US\$140 million, respectively, with a total amount of US\$2.85 billion just in Australia. In the USA alone, US\$6–15 million are spent on chronic wound care annually, while chronic wounds sum 2–3% of the total health expenses in the UK and 2% of the total health expenses in Australia<sup>13–15</sup>. Similar rates can be observed in the European Union (2%) and Scandinavian countries (2–4%)<sup>13</sup>. In addition, 50% of patients with a chronic wound that is not treated properly over a one-year period face serious mortality and morbidity<sup>16</sup>. Furthermore, due to various aetiologies, wound sizes, different treatment protocols and patient population heterogeneity, it is believed these numbers may be underestimated.

A normal inflammatory process ends with cleansing from infectious agents, and total repair of the injured area. In some conditions, however, the immune reaction somehow continues, and the wound becomes persistent. Although the exact mechanisms have not yet been clarified, persistent infection, hypersensitive reactions, long-term exposure to toxic or foreign bodies, and autoimmunity are potential reasons. Studies indicate that malfunctioning immunologic mechanisms, including cytokines and cell stimulants, cause a non-healing wound<sup>17–19</sup>.

The incidence of foot ulcers is rising. Although treatment success rates are promising, relapse rates are very high. Correlating with the rising incidence of diabetes mellitus, diabetic foot ulcers have become a common and important issue that may even lead to amputation. Pressure injuries are another chronic wound type, most prominently observed in long-term hospitalised patients. Studies indicate that in the UK alone, 4% of health expenses are incurred due to pressure injuries<sup>15</sup>. Of note, these wounds are observed usually in high-

risk populations such as disabled and/or elderly persons who are often characterised with major co-morbidities (for example, angiopathic, renal and hepatic problems), and nutritional issues, causing a problematic immune status<sup>20–22</sup>. This condition may result with seriously infected ulcers and also exposure to healthcare-associated or nosocomial infectious agents makes the wound “untreatable” due to probable antimicrobial resistance. To date, *Staphylococcus aureus*, coagulase negative staphylococci (CoNS), *Enterococcus faecalis*, *Proteus* species, anaerobic bacteria, and *Pseudomonas aeruginosa* have the highest isolation rates from chronic wounds<sup>12</sup>. It must be noted that there is great variability between infectious agents<sup>15,23</sup>.

Biofilm formation created by pathogen agents is another issue of chronic wounds. It causes antimicrobial resistance, provokes or aggravates chronic inflammation, prevents healing, and eventually causes treatment failure, or the necessity for long-term, troublesome, and expensive treatments<sup>12</sup>. Biofilm occurs by the aggregation of microorganisms creating a complex multicellular community, resulting in a continuous chain of hyperinflammation, pathogenic invasion of vascular areas, increased capillary permeability and release of the intravascular contents. In addition, biofilm infections are often polymicrobial, which is also another problem to eradicate infection. More than 14 million biofilm infections occur with a mortality of 350,000 individuals, annually<sup>24,25</sup> and biofilm is formed in over 60% of chronic wounds<sup>26</sup>.

There are many types of wound dressings with different specialties designed for various wound types. These dressings include film, hydrogel, foam, hydrocolloid, alginate, hydrofibre, antimicrobial dressings, biologic dressings (MDT), apitherapeutic (honey) products, and tissue engineering products. A good dressing should be easily applicable, aesthetically pleasing, non-painful, cheap, non-allergic, non-toxic, non-traumatising, exude-absorbent, capable of protecting against moisture and warmth, capable of gas transport, preventive of contamination, and capable of necrotic tissue debridement. Aetiologic reason, wound type, and infection status are the main factors considered when choosing the right wound dressing<sup>14,15,23,27</sup>.

Biofilm formation results in a limitation of choices, both for treatment strategies and wound dressings. Guidelines agree on the treatment of biofilms as part of a multi-therapeutic design. Debridement is the major approach, but creating a moisture balance, managing host factors (blood sugar regulation, nutrition, et cetera), eliminating the infectious agents and preventing biofilm re-formation by using topical antiseptics, are all crucial. Debridement alone cannot remove all biofilm and there is only a limited time period before biofilm re-formation occurs. In addition, systemic antimicrobials are usually not effective to degrade biofilms<sup>28</sup>. So, in the treatment of biofilms it is critically important to apply a method(s) that provides strong — easily repeatable — debridement, antimicrobial activity that results in suppressing hyperinflammation.

## MAGGOTS

For use in MDT, a maggot should facilitate debridement by removing necrotic tissue (not living tissue), show antimicrobial and antibiofilm activity, break the immunity chain of the chronic wound, and stimulate wound healing<sup>4</sup>. Many species were investigated for their ability to achieve these goals. Of these, *Lucilia sericata* larvae have been extensively evaluated and found to be effective<sup>4,10</sup>. *Lucilia cuprina* was also reported, especially for diabetic wounds<sup>29</sup>. Many other species such as *Calliphora vicina*, *Calliphora vomitoria*, *Phormia regina*, *Chrysomya albiceps*, *Sarcophaga carnaria*, and *Hermetia illucens* were indicated to have potential for the same use<sup>30</sup>.

The genus *Lucilia* (Diptera: Calliphoridae) consists of over 160 species that have great importance in forensic entomology<sup>31,32</sup>. *Lucilia sericata* (green bottle fly) is common worldwide, but especially in tropical areas, and has a considerable ability to sense carrion from miles away. In a lifetime, a female maggot leaves nearly 200 eggs that quickly grow to their adult forms by passing through various instars and pupal phases. Maggots (larvae) have a complex body form with 12 segments, dark and light sensors, and secretory glands. As they are “dead eaters”, their glands secrete strong proteolytic enzymes, and they can digest nearly half of their body weight in just five minutes. Interestingly, they have a considerable capability to protect themselves from infectious agents, not just in the instar phases, but also in the pupal phase (auto disinfection)<sup>4,10</sup>.

## MAGGOT DEBRIDEMENT THERAPY

Although scientific reports detailing MDT have mostly been published after World War I, MDT has been practised for many years prior to this date<sup>33,34</sup>. In the 1940s, interest in MDT dropped, but it was later revived in the 1990s<sup>5-9</sup>. The effects of MDT on primarily venous stasis ulcers, pressure injuries, neuropathic ulcers (diabetic foot ulcers), traumatic and post-surgical non-healing wounds, burns, arterial ulcers, Buerger disease, cellulitis, mastoiditis, lymphostasis, osteomyelitis and necrotic tumours have been investigated, and the results were very promising<sup>4</sup>. Many studies have been published to date, and recently, MDT has been accorded an important role in chronic wound care in many countries<sup>35-41</sup>.

The species most widely used for this purpose is *Lucilia sericata* larvae<sup>4,10</sup>. Facilities routinely applying MDT produce *L. sericata* in an artificial climate in light/dark condition rooms. At the Instar 2 and 3 stages, maggots are sterilised with various solutions to avoid potential sepsis<sup>36,42-47</sup>. After contamination controls, the maggots are ready to be applied.

MDT can be applied in two forms: “free-range, confinement dressing, cage dressing” or “containment dressing, biobag”. These two methods have their respective advantages. For wounds with deep necrotic tissue, free-range dressing is recommended owing to a higher efficiency debridement and a shorter period for application. However, because of patient

discomfort, complaints (pain, disturbance), and potential rejection for application, many professionals choose to apply biobags<sup>4,38,48</sup>. A few publications indicate the possibility of bleeding due to free-range dressings, but this remains controversial. Maggot movement on the wound, proteolytic reactions, and fibrinolysis may explain this phenomenon. As such, further studies should be performed in this regard<sup>49,50</sup>.

A study investigating MDT from a practitioners’ perspective revealed no difference between these two methods<sup>48</sup>. Furthermore, biobag application may lead to patient cooperation and willingness that may result in increased application numbers<sup>4,38,48</sup>. Although this application may result in higher financial expenses, overall, MDT seems to be much cheaper than conventional wound dressing methods<sup>51</sup>. Evidently, it is too hard to interpret the current data with certainty, and it is recommended that healthcare facilities evaluate the data on an individual basis.

The most problematic issue for MDT is patient acceptance. A few studies have reported that patients may experience pain or report disturbance (restless feelings other than pain)<sup>4</sup>, but we know these factors are strongly variable. Patient acceptance and pain levels may also vary due to wound type. Steenworde *et al.*<sup>52</sup> reported that these patient cooperation issues did not affect the efficiency of MDT unless the patient stopped therapy. They also stated that “acceptance and willingness” are good ways to avoid cooperation issues.

Several contraindications to MDT such as coagulopathies, allergies to larvae/larval secretions, haemorrhagic abscess, and progressive necrotising wounds were reported. In addition, the respiratory system, head area, fistulae to vital organs, endocrine glands, internal organs and open abdominal injuries are not applicable locations for MDT<sup>4,53</sup>.

While there are many studies to highlight the effect mechanisms of MDT, a huge darkness still remains. The mechanisms, to date, can be divided into four titles to facilitate understanding:

### Debridement

The first rule of larval debridement is that the larvae are to be fed necrotic tissue, not living tissue. *Lucilia sericata* larvae employ physical and chemical mechanisms to selectively feed on necrotic tissue. It was found that a maggot can break down 25 mg of necrotic tissue in just 24 hours<sup>54</sup>. Mechanical debridement is achieved by scraping the wound area<sup>55,56</sup>. In fact, maggots search for their food, providing another therapeutic advantage. Even in the deepest areas of the wound, maggots eat necrotic tissue, especially when used in free-range dressing<sup>4,10,38</sup>. The chemical mechanism of debridement is achieved by proteolytic enzymes that are secreted in the digestive system of the maggot. These enzymes also play a key role in the antimicrobial effects of MDT<sup>57,58</sup>. In a recent study, a chymotrypsin-like serine protease, identified from *L. sericata*, affects clotting mechanisms by breaking down extracellular matrix proteins

(fibronectin, laminin and collagen IV), indicating chemical debridement<sup>59</sup>.

#### Antimicrobial effect

Maggots exert their antibacterial effects by both “eating” and their bactericidal activity of excretion and secretion (ES). Mumcuoglu<sup>38</sup> described the mechanical eating activity in 2001. In an academic thesis, Dogandemir<sup>60</sup> investigated both sterile and patient-applied whole-body fluids (WBF) via the microdilution method, and found a greater antibacterial effect exerted by gram-positive bacteria than by gram-negative bacteria. It was also noted that there was no effect on *Pseudomonas aeruginosa* and *Candida albicans*. On the contrary, Margolin et al.<sup>61</sup> reported antifungal activity, but the study was performed with living maggots, not WBFs.

Bexfield et al.<sup>42,43</sup>, Barnes et al.<sup>36</sup>, and Huberman et al.<sup>44,45</sup> studied ES and haemolymph fluids, and found many antibacterial proteins of various molecular masses. Teh et al.<sup>62</sup> studied the antibacterial effects and compounds of larval extracts, and found fatty acids that may have an inhibitory effect on major bacterial growth. Huberman et al.<sup>45</sup> and Kerridge et al.<sup>63</sup> also stated that secretory proteins and their molecular masses vary according to bacterial exposure. They noted that after exposure, early secretory proteins have low molecular masses while late ones have high masses.

Chernysh et al.<sup>64</sup> identified the protein dipterin (8882 and 9025 Da) in maggot ES, and Kruglikova and Chernysh<sup>46</sup> identified new protein molecules with molecular masses of 1014–9025 Da and 174–904 Da. Antibacterial proteins later identified were lucifensin<sup>65,66</sup> and lucifensin II<sup>67</sup>. Interestingly, lucifensin has recently been investigated for its role against biofilm formation, but further research is still required<sup>57</sup>.

Andersen et al.<sup>35</sup> studied antibacterial proteins in maggot ES and their sequences via transposon assisted signal trapping (TAST) technique, and compared their findings using The Basic Local Alignment Search Tool (BLAST) system. Proteins showed 28–91% homology with lectin, defensin, attacin, and kitin binding proteins. They also tested the antimicrobial activity of lucifensin, and found that it had potent activity on gram-positive bacteria, but was less effective against gram-negative bacteria and fungi. Recently, Tellez et al.<sup>68</sup> identified a new protein named Lucilin, a cecropin-like peptide from *Lucilia eximia*, which shows antimicrobial activity especially against gram-negative bacteria and immunomodulatory activity.

In 2014, Valachova et al.<sup>57</sup> revealed the presence in maggot ES of phenyl metalloproteinase, signal peptide protease, three different proteases, and chymotrypsin secretions, stated their amplicons, and, by using BLAST, determined their amino acid sequence homologies with other proteins. In another study, they evaluated larval midgut lysozymes according to molecular mass, amplicons, and antibacterial effects<sup>58</sup>.

Pöppel et al.<sup>69</sup> detected 47 genetic locations from *L. sericata*, and produced 23 synthetic proteins with antifungal activity, including cecropin, cecropin-like, proline rich, stomoxyn, and defensin. Of particular importance are “elevated during infection (edin)” proteins, whose synthesis is upregulated during exposure to infectious agents. Another interesting finding is the observed synergistic and additive effects between proteins. In addition, they noted that these antimicrobial peptides (AMPs) were synthesised in many body parts of the maggot.

Erdmann et al.<sup>70</sup> and Greenberg et al.<sup>71</sup> reported that both phenylacetic acid and phenylacetaldehyde produced by commensal *Proteus mirabilis* in the gastrointestinal system of maggots have antibacterial activity. The same mechanism also led to a bacteriostatic effect by wound surface alkalisation<sup>72</sup>. This alkalisating effect not only results in bacteriostatic activity, but also creates an optimal environment for antimicrobial enzymes<sup>73</sup>.

Pöppel et al.<sup>74</sup> isolated a novel protein molecule, lucimycin, which showed homology with lucifensin. Despite this homology, lucimycin, unlike lucifensin, showed antifungal activity. Although these researchers studied a limited number of fungi, they observed inhibitory effects on spore germination and mycelial growth for particular species (*Cladosporium herbarium*, *Alternaria alternata*, *Lichtheimia corymbifera*, *Mucor circinelloides*, *Candida albicans*, and *Trichosporon asahii*). In contrast, no antifungal activity was observed against some species (*Rhizopus oryzae*, *Aspergillus fumigatus*, *Aspergillus terreus*, *Arthoderma benhamiae*).

Another study, published by Polat et al.<sup>75</sup>, led to a new perspective on MDT. They investigated, *in vivo*, the anti-leishmanial effects of *L. sericata* ES on *Leishmania tropica* infected mice. They reported that fresh and pure ES significantly advanced lesion healing, but this anti-leishmanial effect could not be observed on promastigotes from culture. They suggested that quick oxidation of ES might be the reason for this discrepancy. Daeschlein et al.<sup>76</sup> reported that the antibacterial activity of ES against *E. coli* and *S. aureus* decreased over time, which supports this quick destabilisation theory. Sanei-Dehkordi et al.<sup>77</sup> supported this data in a study on *Leishmania major*. They found that treatment with *L. sericata* ES led to a considerably lower number of amastigote-infected macrophages in a test group than that in the control group *in vitro*, and observed significant changes in lesion size, indicating that ES inhibits leishmanial virulence.

The common perception is that maggots are more effective against gram-positive bacteria, and that the antibacterial effects on *Pseudomonas* and *Acinetobacter* species are somehow more limited than on other species<sup>35,78,79</sup>. Antifungal and anti-leishmanial effects have also been observed<sup>74,75</sup>, but must still be supported by further investigation.

### Effects on wound healing

Several studies have reported that MDT promotes wound healing<sup>80-83</sup>. As previously stated, this may be due to both the cleansing of necrotic tissue and antimicrobial effects. However, many researchers reported direct activities of larvae on wound healing. Van der Plas *et al.*<sup>84</sup> studied the cellular effects of *L. sericata* ES, and found that it has inhibitory effects on the proinflammatory immune response (which is a persistent problem in chronic inflammation) without any negative impact on the antimicrobial effects of neutrophils. In addition, decreases in the level of elastase secretion, neutrophil chemotaxis, and hydrogen peroxide generation were observed in a dose-dependent fashion. In contrast, neutrophil chemotaxis and antimicrobial effects were not affected. Another study reported that monocyte proinflammatory and cytokine responses were considerably affected by ES<sup>85</sup>. Tamura *et al.*<sup>86</sup> reported that ES had inhibitory effects on the complement system, resulting in the suppression of persistent inflammation without any negative impact on neutrophil antimicrobial action. It has also been found that proteolytic ES activity has a positive impact on extracellular matrix mechanisms, which also play a role in wound healing<sup>87</sup>. Baumann *et al.*<sup>88</sup> uncovered urate oxidase and allantoin mechanisms in *L. sericata* larvae that regulate pH levels in the wound. This is also needed to ensure the potency of antimicrobial effects and the enzymatic wound healing processes.

Neoangiogenesis is an important part of wound healing, and beneficial effects of *L. sericata* ES on neoangiogenesis have been reported<sup>89</sup>. In contrast, Singorenko *et al.*<sup>90</sup> reported that ES had no direct effect on wound healing. Although they demonstrated transcriptional changes, no positive impact on cell viability, proliferation, migration, or angiogenesis was observed. However, they revealed significant effects on the immune response, which may indirectly modulate wound healing. A recent study has, however, reported that MDT stimulates endothelial cell proliferation and neoangiogenesis in diabetic foot wounds<sup>91</sup>. Zhang *et al.*<sup>92</sup> reported strong evidence that neoangiogenesis was actually stimulated by MDT. They demonstrated a remarkable increase in miR-126 expression (an mRNA that stimulates neoangiogenesis) in MDT-treated patients with diabetic foot ulcers, and confirmed these findings by observing increased miR-126 expression *in vitro* in human umbilical vein endothelial cells. Methodological difference could be a factor in these conflicting results. In addition, wound type and aetiology may have an impact on MDT effects.

### Biofilm degradation effect

Biofilm formation is a serious problem that causes antimicrobial resistance, added medical expenses, and additional long-term treatment<sup>93</sup>. Cazander *et al.* published two different studies<sup>37,94</sup>, indicating that ES showed a degrading effect on *S. aureus*, *S. epidermidis*, *K. oxytoca*, *E. faecalis*, *E. cloacae*, and *P. aeruginosa* biofilms on polyethylene, titanium, and surgical steel surfaces. Van der

Plas *et al.*<sup>95</sup> found that ES broke down *S. aureus* biofilm without any additional antibiotic treatment. They also noted that linezolid and clindamycin could not degrade biofilm in the first 24 hours.

*Pseudomonas aeruginosa* causes problems, especially in intensive care units, due to both its ability to create biofilms, and its multidrug resistance<sup>12</sup>. Although the anti-pseudomonal effect of MDT is currently controversial, Cazander *et al.*<sup>37,94</sup> observed a degrading effect of ES on *P. aeruginosa* biofilms, and Pöppel *et al.*<sup>69</sup> indicated genetic markers for antimicrobial effects against *P. aeruginosa*. Brown *et al.*<sup>96</sup> identified DNAase-1 in *L. sericata* ES, which had a destructive effect on *P. aeruginosa* biofilms.

### FUTURE PERSPECTIVE

MDT itself is in use of clinical practice; however, researchers can be divided into two sections about this subject: i) those who practise MDT in their studies in favour of clinical perspective; and ii) those who perform studies in a molecular and proteomic vision. These two “cults” actually walk on different paths to the same target. Although MDT has a great scientific value in clinical practice due to serial studies reported worldwide, only a few researchers focused at the molecular and protein levels. Recently, several studies published have especially chosen the subject area of specific molecules isolated from medicinal maggots and their potential beneficial effects as antimicrobials, wound healers, and so on. Recent studies of Pöppel *et al.*<sup>59,69</sup> showed recombinant technology can be a tool for gaining particular molecules singularly to use for medicinal purposes, which also opened a path to transgenic molecules<sup>97</sup>. In another study, Gordya *et al.*<sup>98</sup> reported anti-biofilm activity of *Calliphora vicina* AMPs by significantly destroying the matrix and eliminating the bacteria. These data may lead medical societies to produce new wound dressings containing components such as these particular AMPs and transgenic molecules for especially the occasions such as biofilms, where the clinicians have limited options. Despite it being hard to obtain high amounts of purified proteins directly from maggots, thanks to recombinant science, these molecules are still strong candidates for future drug researches<sup>99</sup>.

### CONCLUSIONS

As a complementary medicinal technique, MDT stands as a “scientifically proven” method. Although there are still unclarified issues, the use of MDT is undisputed. Chronic wounds are a common problem and MDT acts as another therapeutic option for patients.

Antibacterial activity against gram-positive bacteria is clearly observed, but effects on other particular bacteria (*Acinetobacter* spp and *Pseudomonas aeruginosa*) and fungi remain still controversial<sup>35,78,79</sup>. Previously stated studies indicate many mechanisms and molecules in action, but it seems these are only single drops in a vast ocean. Recently, Franta *et al.*<sup>100</sup> found many secreted enzymes of *L. sericata*

that have the potential to play a role in several effect mechanisms. It should be noted that these mechanisms and molecules do not just work alone, but work synergistically. In addition, it seems there is a “provocation and adaptation” mechanism in which peptides are selectively secreted, depending on pathogen exposure or peptide type (molecular mass and activity), and the maggot somehow “adapts” to what it encounters. Some studies indicate that the exact antimicrobial effect comes forward in the situation of this so-called “provocation and adaptation,” and observation of the effects on these controversial species may be achieved in this period. In addition, transgenic maggots may be more cost-effective and improve patient outcomes by secreting particular wound-healing peptides<sup>97</sup>. Further investigations are necessary.

Chronic wounds have many aetiologic reasons that affect the success rate of MDT. Most reasons are systemic diseases (for example, diabetes mellitus) with co-morbidities (for example, chronic renal failure) that also negatively impact the immune system of the patient. This situation strongly limits the applicability of systemic drugs, and forces medical professionals to avoid the possible side effects. In addition, chronic wounds such as venous stasis and diabetic foot ulcerations actually have a vascularisation pathology, which may also limit the benefits of systemic treatments. MDT has another advantage in that it is applied locally, and has no systemic effect. This makes MDT especially appealing for patients with systemic co-morbidities<sup>101</sup>.

In conclusion, MDT satisfies nearly all expectations for optimal wound care: fighting infection, debridement, provoking wound healing, and neoangiogenesis<sup>4,89</sup>. Its low-cost, noninvasive nature, and absence of systemic effects make this method a good option. MDT is not an exact wound treatment method, but can be an important part of a multidisciplinary approach to treat chronic wounds. There is a black hole in its mechanisms, and by highlighting them, the area of use might widen.

## CONFLICT OF INTEREST

The authors declare no conflict of interests.

## FUNDING

This article did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## ACKNOWLEDGEMENTS

We declare special thanks to Prof Sibel Erguven, PhD (Hacettepe University, Faculty of Medicine, Department of Medical Microbiology, Ankara, Turkey) for her precious guidance.

## REFERENCES

- Fischer FH, Lewith G, Witt CM *et al*. High prevalence but limited evidence in complementary and alternative medicine: guidelines for future research. *BMC Complement Altern Med* 2014;14(1): 46.
- World Health Organization (WHO). WHO traditional medicine strategy: 2014–2023. Geneva, Switzerland: WHO, 2013.
- Barnes PM, Bloom B, Nahin RL. Complementary and alternative medicine use among adults and children: United States, 2007. *National health statistics reports*; No. 12. Hyattsville, MD: National Center for Health Statistics, 2008.
- Sherman RA, Mumcuoglu KY, Grassberger M, Tantawi TI. Maggot Therapy. In: Grassberger M, Sherman RA, Gileva OS, Kim CMH, Mumcuoglu KY (eds). *Biotherapy — history, principles and practice: A practical guide to the diagnosis and treatment of disease using living organisms*. Amsterdam: Springer Science & Business Media, 2013, pp. 5–29.
- Sherman RA, Hall M, Thomas S. Medicinal maggots: an ancient remedy for some contemporary afflictions. *Annu Rev Entomol* 2000;45(1):55–81.
- Sherman RA, Pechter EA. Maggot therapy: a review of the therapeutic applications of fly larvae in human medicine, especially for treating osteomyelitis. *Med Vet Entomol* 1988;2(3):225–230.
- Stoddard S, Sherman R, Mason B, Pelsang D, Sherman R. Maggot debridement therapy. An alternative treatment for nonhealing ulcers. *J Am Podiatr Med Assoc* 1995;85(4):218–221.
- Sherman RA, Tran JMT, Sullivan R. Maggot therapy for venous stasis ulcers. *Arch Dermatol* 1996;132(3):254–256.
- Sherman RA, Wyle FA. Low-cost, low-maintenance rearing of maggots in hospitals, clinics, and schools. *Am J Trop Med Hyg* 1996;54(1):38–41.
- Fleischmann W, Grassberger M, Sherman RA. *Maggot therapy: A handbook of maggot-assisted wound healing*. London: Thieme Publications, 2004.
- Game FL, Apelqvist J, Attinger C *et al*. IWGDF Guidance on use of interventions to enhance healing of chronic ulcers of the foot in diabetes. *International Working Group on Wound Healing Document*, Amsterdam, 2015.
- Trøstrup H, Bjarnsholt T, Kirketerp-Møller K, Høiby N, Moser C. What is new in the understanding of non healing wounds epidemiology, pathophysiology and therapies. *Ulcers* 2013;8:1–6.
- Graves N, Zheng H. Modelling the direct health care costs of chronic wounds in Australia. *Wound Pract Res* 2014;22(1):20–33.
- Sood A, Granick MS, Tomaselli NL. Wound dressings and comparative effectiveness data. *Adv Wound Care* 2014;3(8):511–529.
- Nunan R, Harding KG, Martin P. Clinical challenges of chronic wounds: searching for an optimal animal model to recapitulate their complexity. *Dis Model Mech* 2014;7(11):1205–1213.
- Cazander G, Jukema GN, Nibbering PH. Complement activation and inhibition in wound healing. *Clin Dev Immunol* 2012;8:1–14.
- Kumar V, Abbas AK, Aster JC. *Robbins and Cotran pathologic basis of disease*, 9th edn. London: Elsevier Publications, 2015.
- Nathan C, Ding A. Nonresolving inflammation. *Cell* 2010;140(6):871–882.
- Van der Plas MJA. Effect of maggot secretions on microbiological, haematological and immunological processes. *Maggot Therapy's Modes of Action*. EZ Leiden, Leiden University Publication 2009, pp. 9–35.
- Gorecki C, Brown JM, Nelson EA *et al*. Impact of pressure ulcers on quality of life in older patients: a systematic review. *J Am Geriatr Soc* 2009;57(7):1175–1183.
- Reddy M, Gill SS, Rochon PA. Preventing pressure ulcers: a systematic review. *JAMA* 2006;296(8):974–984.
- Banks M, Bauer J, Graves N, Ash S. Malnutrition and pressure ulcer risk in adults in Australian health care facilities. *Nutrition* 2010;26(9):896–901.
- Dreifke MB, Jayasuriya AA, Jayasuriya AC. Current wound healing procedures and potential care. *Mater Sci Eng C* 2015;48:651–662.

24. Percival SL, Hill KE, Williams DW, Hooper SJ, Thomas DW, Costerton JW. A review of the scientific evidence for biofilms in wounds. *Wound Repair Regen* 2012;20(5):647–657.
25. Wolcott RD, Rhoads DD, Bennett ME et al. Chronic wounds and the medical biofilm paradigm. *J Wound Care* 2010;19(2):45–53.
26. Malone M, Bjarnsholt T, McBain AJ et al. The prevalence of biofilms in chronic wounds: a systematic review and meta-analysis of published data. *J Wound Care* 2017;26(1):20–25.
27. Broussard KC, Powers JG. Wound dressings: selecting the most appropriate type. *Am J Clin Dermatol* 2013;14(6):449–459.
28. Schultz G, Bjarnsholt T, James GA et al. Consensus guidelines for the identification and treatment of biofilms in chronic nonhealing wounds. *Wound Repair Regen* 2017;25(5):744–757.
29. Paul AG, Ahmad NW, Lee HL et al. Maggot debridement therapy with *Lucilia cuprina*: a comparison with conventional debridement in diabetic foot ulcers. *Int Wound J* 2009;6:39–46.
30. Raposio E, Bortolini S, Maistrello L, Grasso DA. Larval therapy for chronic cutaneous ulcers: Historical review and future perspectives. *Wounds* 2017;29(12):367–373.
31. Stevens J, Wall R. Classification of the genus *Lucilia* (Diptera: Calliphoridae): A preliminary parsimony analysis. *J Nat Hist* 1996;30(7):1087–1094.
32. Wells JD, Wall R, Stevens JR. Phylogenetic analysis of forensically important *Lucilia* flies based on cytochrome oxidase I sequence: a cautionary tale for forensic species determination. *Int J Legal Med* 2007;121(3):229–233.
33. Baer WS. Sacro-iliac joint — Arthritis deformans — viable antiseptic in chronic osteomyelitis. *Proc Int Assembly Interstate Postgrad Med Assoc North Am* 1929;371:365–372.
34. Baer WS. The treatment of chronic osteomyelitis with the maggot (larva of the blow fly). *J Bone Joint Surg* 1931;13(3):438–475.
35. Andersen AS, Sandvang D, Schnorr KM et al. A novel approach to the antimicrobial activity of maggot debridement therapy. *J Antimicrob Chemother* 2010;65(8):1646–1654.
36. Barnes KM, Dixon RA, Gennard DE. The antibacterial potency of the medicinal maggot, *Lucilia sericata* (Meigen): variation in laboratory evaluation. *J Microbiol Methods* 2010;82(3):234–237.
37. Cazander G, Van de Veerdonk MC, Vandenbroucke-Graults CM, Schreurs MW, Jukema GN. Maggot excretions inhibit biofilm formation on biomaterials. *Clin Orthop Relat Res* 2010;468(10):2789–2796.
38. Mumcuoglu KY. Clinical applications for maggots in wound care. *Am J Clin Dermatol* 2001;2(4):219–227.
39. Whitaker IS, Twine C, Whitaker MJ, Welck M, Brown CS, Shandall A. Larval therapy from antiquity to the present day: mechanisms of action, clinical applications and future potential. *Postgrad Med J* 2007;83(980):409–143.
40. Tanyuksel M, Araz E, Dundar K et al. Maggot debridement therapy in the treatment of chronic wounds in a military hospital setup in Turkey. *Dermatol* 2004;210(2):115–118.
41. Polat E, Cakan H, Ipek T. Larva debridement therapy (LDT). *Turkish Journal of Family Practice* 2011;14(4):188–191.
42. Bexfield A, Bond AE, Roberts EC et al. The antibacterial activity against MRSA strains and other bacteria of a < 500 Da fraction from maggot excretions/secretions of *Lucilia sericata* (Diptera: Calliphoridae). *Microbes Infect* 2008;10(4):325–333.
43. Bexfield A, Nigam Y, Thomas S, Ratcliffe NA. Detection and partial characterisation of two antibacterial factors from the excretions/secretions of the medicinal maggot *Lucilia sericata* and their activity against methicillin-resistant *Staphylococcus aureus* (MRSA). *Microbes Infect* 2004;6(14):1297–1304.
44. Huberman L, Gollop N, Mumcuoglu KY, Block C, Galun R. Antibacterial properties of whole body extracts and haemolymph of *Lucilia sericata* maggots. *J Wound Care* 2007;16(3):123–127.
45. Huberman L, Gollop N, Mumcuoglu KY et al. Antibacterial substances of low molecular weight isolated from the blowfly, *Lucilia sericata*. *Med Vet Entomol* 2007;21(2):127–131.
46. Kruglikova A, Chernysh S. Antimicrobial compounds from the excretions of surgical maggots, *Lucilia sericata* (Meigen) (Diptera, Calliphoridae). *Entomological Review* 2011;91(7):813–819.
47. Limsopatham K, Khamnoi P, Sukontason KL, Boonyawan D, Chaiwong T, Sukontason K. Sterilization of blow fly eggs, *Chrysomya megacephala* and *Lucilia cuprina* (Diptera: Calliphoridae) for maggot debridement therapy application. *Parasitol Res* 2017; pp 1–9. doi:10.1007/s00436-017-5435-9
48. Mudge E, Price P, Neal W, Harding KG. A randomized controlled trial of larval therapy for the debridement of leg ulcers: Results of a multicenter, randomized, controlled, open, observer blind, parallel group study. *Wound Repair Regen* 2014;22(1):43–51.
49. Church J. Re: Bleeding complications in patients treated with maggot debridement therapy. *Steenvoorde P & Oskam J. IJLEW* 2005;4(1):57–58. *Int J Low Extrem Wounds* 2005;4(1):59.
50. Steenvoorde P, Jacobi CE, Oskam J. Maggot debridement therapy: free-range or contained? An *in-vivo* study. *Adv Skin Wound Care* 2005;18(8):430–435.
51. Thomas S. Cost of managing chronic wounds in the UK, with particular emphasis on maggot debridement therapy. *J Wound Care* 2006;15(10):465–469.
52. Steenvoorde P, Buddingh TJ, Engeland AV, Oskam J. Maggot therapy and the “Yuk” factor: An issue for the patient? *Wound Repair Regen* 2005;13(3):350–352.
53. Jordan A, Khiyani N, Bowers SR, Lukaszczuk JJ, Stawicki SP. Maggot debridement therapy: A practical review. *Int J Acad Med* 2018;4(1):21–34.
54. Sherman RA. Maggot therapy for foot and leg wounds. *Int J Low Extrem Wounds* 2002;1(2):135–142.
55. Barnard D. Skeletal-muscular mechanisms of the larva of *Lucilia sericata* (Meigen) in relation to feeding habit (Diptera: Calliphoridae). *Pan-Pac Entomol* 1977;3(1):34–38.
56. Thomas S, Wynn K, Fowler T, Jones M. The effect of containment on the properties of sterile maggots. *Br J Nurs* 2002;11(Supp2):21–28.
57. Valachova I, Majtan T, Takac P, Majtan J. Identification and characterisation of different proteases in *Lucilia sericata* medicinal maggots involved in maggot debridement therapy. *J Appl Biomed* 2014;12(3):171–177.
58. Valachova I, Takac P, Majtan J. Midgut lysozymes of *Lucilia sericata* — new antimicrobials involved in maggot debridement therapy. *Insect Mol Biol* 2014;23(6):779–787.
59. Pöppel AK, Kahl M, Baumann A, Wiesner J, Gokcen A, Beckert A. A Jonah-like chymotrypsin from the therapeutic maggot *Lucilia sericata* plays a role in wound debridement and coagulation. *Insect Biochem Mol Biol* 2016;70:138–147.
60. Dogandemir G. Investigation of antimicrobial activities of *Lucilia sericata* against microorganisms colonising on chronic wounds. *Medical Microbiology Thesis, Gulhane Military Medical Academy, Ankara, Turkey, 2010.*
61. Margolin L, Gialanella P. Assessment of the antimicrobial properties of maggots. *Int Wound J* 2010;7(3):202–204.
62. Teh CH, Nazni WA, Norazah A, Lee HL. Determination of antibacterial activity and minimum inhibitory concentration of larval extract of fly via resazurin-based turbidometric assay. *BMC Microbiology* 2017;17(1):36.
63. Kerridge A, Lappin-Scott H, Stevens J. Antibacterial properties of larval secretions of the blowfly, *Lucilia sericata*. *Med Vet Entomol* 2005;19(3):333–337.
64. Chernysh SI, Gordja NA, Simonenko NP. Diapause and immune response: induction of antimicrobial peptides synthesis in the blowfly, *Calliphora vicina* R.-D. (Diptera: Calliphoridae). *J Entomol Sci* 2000;3(1):139–144.
65. Čeřovský V, Žďárek J, Fučík V, Monincová L, Voburka Z, Bém R. Lucifensin, the long-sought antimicrobial factor of medicinal maggots of the blowfly *Lucilia sericata*. *Cell Mol Life Sci* 2010;67(3):455–466.

66. Valachová I, Bohová J, Pálošová Z, Takáč P, Kozánek M, Majtán J. Expression of lucifensin in *Lucilia sericata* medicinal maggots in infected environments. *Cell Tissue Res* 2013;353(1):165–171.
67. El Shazely B, Veverka V, Fučík V, Voburka Z, Žďárek J, Čeřovský V. Lucifensin II, a defensin of medicinal maggots of the blowfly *Lucilia cuprina* (Diptera: Calliphoridae). *J Med Entomol* 2013;50(3):571–578.
68. Tellez GA, Zapata JA, Toro LJ et al. Identification, characterization, immunolocalization, and biological activity of Lucilin peptide. *Acta Trop* 2018;185:318–326.
69. Pöppel AK, Vogel H, Wiesner J, Vilcinskas A. Antimicrobial peptides expressed in medicinal maggots of the blow fly *Lucilia sericata* show combinatorial activity against bacteria. *Antimicrob Agents Chemother* 2015;59(5):2508–2514.
70. Erdmann G, Khalil S. Isolation and identification of two antibacterial agents produced by a strain of *Proteus mirabilis* isolated from larvae of the screwworm (*Cochliomyia hominivorax*) (Diptera: Calliphoridae). *J Med Entomol* 1986;23(2):208–211.
71. Greenberg B. Model for destruction of bacteria in the midgut of blow fly maggots. *J Med Entomol* 1968;5(1):31–38.
72. Robinson W, Baker F. The enzyme urease and the occurrence of ammonia in maggot-infected wounds. *J Parasitol* 1939;25(2):149–155.
73. Chambers L, Woodrow S, Brown AP et al. Degradation of extracellular matrix components by defined proteinases from the greenbottle larva *Lucilia sericata* used for the clinical debridement of non-healing wounds. *Br J Dermatol* 2003;148(1):14–23.
74. Pöppel AK, Koch A, Kogel KH et al. Lucimycin, an antifungal peptide from the therapeutic maggot of the common green bottle fly *Lucilia sericata*. *Biol Chem* 2014;395(6):649–656.
75. Polat E, Cakan H, Aslan M et al. Detection of anti-leishmanial effect of the *Lucilia sericata* larval secretions *in vitro* and *in vivo* on *Leishmania tropica*: first work. *Exp Parasitol* 2012;132(2):129–134.
76. Daeschlein G, Mumcuoglu KY, Assadian O, Hoffmeister B, Kramer A. *In vitro* antibacterial activity of *Lucilia sericata* maggot secretions. *Skin Pharmacol Physiol* 2006;20(2):112–115.
77. Sanei-Dehkordi A, Khamesipour A, Akbarzadeh K et al. Anti *Leishmania* activity of *Lucilia sericata* and *Calliphora vicina* maggots in laboratory models. *Experimental Parasitology* 2016;170:59–65.
78. Čeřovský V, Bém R. Lucifensins, the insect defensins of biomedical importance: the story behind maggot therapy. *Pharmaceuticals* 2014;7(3):251–264.
79. Jaklič D, Lapanje A, Zupančič K, Smrke D, Gunde-Cimerman N. Selective antimicrobial activity of maggots against pathogenic bacteria. *J Med Microbiol* 2008;57(5):617–625.
80. Armstrong DG, Salas P, Short B et al. Maggot therapy in “lower-extremity hospice” wound care: fewer amputations and more antibiotic-free days. *J Am Podiatr Med Assoc* 2005;95(3):254–257.
81. Dumville JC, Worthy G, Soares MO et al. VenUS II: a randomised controlled trial of larval therapy in the management of leg ulcers. *Health Technol Assess* 2009;1–220.
82. Sherman RA. Maggot versus conservative debridement therapy for the treatment of pressure ulcers. *Wound Repair Regen* 2002;10(4):208–214.
83. Sherman RA. Maggot therapy for treating diabetic foot ulcers unresponsive to conventional therapy. *Diabetes Care* 2003;26(2):446–451.
84. Van der Plas MJ, van der Does AM, Baldry M et al. Maggot excretions/secretions inhibit multiple neutrophil pro-inflammatory responses. *Microbes Infect* 2007;9(4):507–514.
85. Van der Plas MJ, Baldry M, van Dissel JT, Jukema GN, Nibbering PH. Maggot secretions suppress pro-inflammatory responses of human monocytes through elevation of cyclic AMP. *Diabetologia* 2009;52(9):1962–1970.
86. Tamura T, Cazander G, Rooijackers SH, Trouw LA, Nibbering PH. Excretions/secretions from medicinal larvae (*Lucilia sericata*) inhibit complement activation by two mechanisms. *Wound Repair Regen* 2017;25(1):41–50.
87. Horobin AJ, Shakesheff KM, Woodrow S, Robinson C, Pritchard DI. Maggots and wound healing: an investigation of the effects of secretions from *Lucilia sericata* larvae upon interactions between human dermal fibroblasts and extracellular matrix components. *Br J Dermatol* 2003;148(5):923–933.
88. Baumann A, Skaljic M, Lehmann R, Vilcinskas A, Franta Z. Urate oxidase produced by *Lucilia sericata* medical maggots is localized in Malpighian tubes and facilitates allantoin production. *Insect Biochem Mol Biol* 2017;83:44–53.
89. Bexfield A, Bond AE, Morgan C et al. Amino acid derivatives from *Lucilia sericata* excretions/secretions may contribute to the beneficial effects of maggot therapy via increased angiogenesis. *Br J Dermatol* 2010;162(3):554–562.
90. Daurus Singorenko P, Rosario R, Windsor JA, Phillips AR, Blenkiron C. The transcriptional responses of cultured wound cells to the excretions and secretions of medicinal *Lucilia sericata* larvae. *Wound Repair Regen* 2017;25(1):51–61.
91. Sun X, Chen JA, Zhang J, Wang W, Sun J, Wang A. Maggot debridement therapy promotes diabetic foot wound healing by up-regulating endothelial cell activity. *J Diabetes Complications* 2016;30(2):318–322.
92. Zhang J, Sun XJ, Chen JA et al. Increasing the miR-126 expression in the peripheral blood of patients with diabetic foot ulcers treated with maggot debridement therapy. *J Diabetes Complications* 2017;31(1):241–244.
93. Römling U, Balsalobre C. Biofilm infections, their resilience to therapy and innovative treatment strategies. *J Intern Med* 2012;272(6):541–561.
94. Cazander G, van Veen KE, Bouwman LH, Bernards AT, Jukema GN. The influence of maggot excretions on PAO1 biofilm formation on different biomaterials. *Clin Orthop Relat Res* 2009;467(2):536–545.
95. Van Der Plas MJ, Jukema GN, Wai SW et al. Maggot excretions/secretions are differentially effective against biofilms of *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 2008;61(1):117–122.
96. Brown A, Horobin A, Blount DG et al. Blow fly *Lucilia sericata* nuclease digests DNA associated with wound slough/eschar and with *Pseudomonas aeruginosa* biofilm. *Med Vet Entomol* 2012;26(4):432–439.
97. Linger RJ, Belikoff EJ, Yan Y et al. Towards next generation maggot debridement therapy: transgenic *Lucilia sericata* larvae that produce and secrete a human growth factor. *BMC Biotechnol* 2016;16(1):30.
98. Gordya N, Yakovlev A, Kruglikova A et al. Natural antimicrobial peptide complexes in the fighting of antibiotic resistant biofilms: *Calliphora vicina* medicinal maggots. *PLoS ONE* 2017;12(3):e0173559.
99. Yan L, Chu J, Li M et al. Pharmacological Properties of the Medical Maggot: A Novel Therapy Overview. *Evid Based Complement Alternat Med* 2018;4934890. doi: 10.1155/2018/4934890
100. Franta Z, Vogel H, Lehmann R, Rupp O, Goesmann A, Vilcinskas A. Next generation sequencing identifies five major classes of potentially therapeutic enzymes secreted by *Lucilia sericata* medical maggots. *Biomed Res Int* 2016;8285428. doi: 10.1155/2016/8285428
101. Cherniack EP. Bugs as drugs, Part 1: Insects: the “new” alternative medicine for the 21st century. *Altern Med Rev* 2010;15(2):124–135.