

Evidence Summary: Wound management-low resource communities: citric acid as a topical antiseptic

May 2016

Author

Wound Healing and Management Node Group — R Watts, E Frehner

QUESTION

What is the best available evidence regarding the use of citric acid for wound management?

CLINICAL BOTTOM LINE

Citric acid has been used as a low-cost option for managing a wide range of chronic wounds which have not responded to conventional management. Sensitivity of a range of bacteria to 3% citric acid, including multiple antibiotic-resistant organisms, has been demonstrated through microscopy, culture and sensitivity (MC&S) (Level 5c evidence). Complete wound healing has been achieved in a high percentage of cases with daily application of 3% citric acid (Levels 2c, 4b, 4c, 4d evidence) and histological findings suggest that citric acid may not create an environment that is toxic to a healing wound (Level 5c evidence). Some patients report mild to moderate irritation for several minutes after application (Level 4b, 4c, 4d evidence, although with superficial burns and large raw areas the burning sensation can be severe (Level 2c evidence). Randomised controlled trials on the effectiveness and safety of citric acid as an antiseptic with large sample sizes are required.

Types of wounds

Evidence on the use of citric acid for wound irrigation and/or topical application has been reported for a range of wounds including burns,¹ foot ulcers,^{2,3} traumatic wounds,⁴ necrotising fasciitis,⁵ bursitis,⁶ lepromatous (Hansen's disease) ulcers⁷ and surgical site infections.⁸⁻¹⁰

Microbiology and histology

A study tested the effectiveness of four topical antiseptics against multi-drug resistant *Pseudomonas aeruginosa* in full thickness burns in rats. Only two of the antiseptic dressings — nanocrystalline silver-coated and silver sulphadiazine 1% dressings — were effective ($p < 0.05$), while the results for chlorhexidine acetate and citric acid 3% were not significant.¹¹ (Level 5c)

In a study of patients with chronic wounds infected with multiple antibiotic-resistant *Escherichia coli*, the bacteria were eliminated and the wounds completely healed in all 34 patients following daily application of 3% citric acid (range 7–42), with diabetic foot and Hansen's disease ulcers taking the longest to heal.³ (Level 4b)

In two cases M&CS conducted before and after application of citric acid to non-diabetic foot ulcers showed eradication of multiple antibiotic-resistant *Staphylococcus aureus* after six treatments and *E. coli* after two treatments. The number of colonies cultured before the application of citric acid was unclear, although in the second case there was active pus

discharge (i.e. the ulcer in the first case may not have been clinically infected) Histological findings from the same study showed development of blood vessels and proliferating fibroblasts after two to six days in the ulcers irrigated daily with 3% citric acid and dressed with a 3% citric acid soaked pad.¹² (Level 4c evidence)

In cases in which MC&S had been performed, other organisms that were sensitive to 3% citric acid included streptococci, *Klebsiella* spp, *Citrobacter* spp, *Staphylococcus albus* and *Proteus* spp. In a number of cases the isolated organisms were found to be sensitive to citric acid while being resistant to a wide range of antibiotics.^{1,2,4,7,8,13} (Level 5c evidence)

The inhibition of bacterial growth is thought to be due to citric acid's reduction of the pH of the wound.^{2,14} The minimum inhibitory concentration (MIC) ranged from 500 to 2500 µg/ml (the MIC being the lowest concentration to which citric acid can be diluted and still remain effective).^{7,8} (Level 5c evidence)

Promising results have been obtained in *in-vivo* studies of a citric acid-based, biodegradable hydrogel dressing material in terms of antimicrobial effects and compatibility with human fibroblasts.¹⁵

Effectiveness in promoting healing

A small double blinded, randomised control trial (n= 50) compared the effectiveness of daily applications of 3% citric acid and Eusol in healing lower limb, muscle deep ulcers resulting from trauma more than 10 days before presentation. (NB. Eusol should no longer be used for any wound management.) The mean time to development of healthy granulation tissue in the citric acid group was 10.56 days compared to 20.04 days in the group treated with Eusol ($p=0.00$). The biopsied tissues from the citric acid group showed a substantial concentration of fibroblasts and neovascularisation at day 7 while these were sparse in the Eusol group tissues. The group treated with citric acid dressings consequently had much shorter hospital stays for approximately half the cost when compared to the group treated with Eusol. Forty percent of the participants were diabetics divided almost equally between the two groups. When compared with the non-diabetic patients in their group, the mean length of hospital stay for the diabetics in the citric acid group was almost the same as the non-diabetics (16.38 days vs 15.78 days). In the Eusol group the mean length of stay for the diabetics was 31 days compared to 24.04 days for non-diabetics in this group.¹⁴

A quasi-experimental study was undertaken with 66 participants with clinically infected burns that had not responded to conventional treatment. In the intervention group (n=46, 5–60% superficial to deep burns) daily application of 3%

citric acid in paraffin was associated with complete healing in 40 (87%) cases in 7–25 days, while healthy granulation tissue formed in the remaining six patients. There was no significant statistical difference ($p=0.145$) in overall healing compared with the control group ($n=20$, 10–70% superficial to deep burns) whose burns were treated with daily topical silver sulfadiazine and systemic antibiotics — 9 cases (45%). Participants in the antibiotics group had more severe burns in terms of depth and total body area coverage.¹ (Level 2c evidence)

In a large prospective observational study ($N= 259$) of non-diabetic patients with traumatic wounds which had not responded to antibiotics and daily povidone iodine dressings, 244 (94%) wounds healed completely with debridement followed by 5–25 days of daily normal saline irrigation and application of 3% citric acid (the remaining six patients required skin grafting). A significant reduction in exudate and pain was observed after 2–3 applications of citric acid.⁴ (Level 4b)

In another uncontrolled observational study ($N=115$) of patients with clinically infected diabetic foot ulcers (Wagner grades I–III) not responding to conventional treatment, 94% ($n=32$) of grade I ulcers treated daily with 3% citric acid in paraffin healed in 5 to 30 days. Ninety four percent ($n=49$) of Wagner grade II ulcers and 86% ($n=25$) of Wagner grade III healed in 16 to 34 days. The majority of ulcers that did not completely heal developed healthy granulation and skin grafts were performed (Level 4b evidence). (Note: patients with

peripheral vascular disease were excluded from the study.)² (Level 4b)

Similar results were obtained in a study of 70 patients with surgical site infections that were not responding to conventional treatment. Daily application of 3% citric acid ointment for 6–25 days resulted in complete healing or formation of healthy granulating tissue allowing final suturing of gaps in the wound in all but one case.⁸ (Level 4b)

In a study of 34 patients with Hansen's disease and ulcers that were not responding to antibiotic therapy, daily applications of 3% citric acid eliminated the pathogens and completely healed all the ulcers in 25 patients (74%). In the remaining nine patients healthy granulation tissue formed but complete healing did not occur, possibly due to underlying nerve and blood vessel damage from the disease.⁷ (Level 4b)

Another small study trialed the use of citric acid to prepare the wound bed for grafting in five road trauma cases with large raw areas infected with multiple antibiotic-resistant bacteria. Healthy granulation tissue was generated in four patients (80%) but deep seated infection could not be eliminated in the fifth patient.¹³ (Level 4b)

In another ten case studies, a variety of clinically infected chronic wounds treated with 3% citric acid achieved complete healing ($n=8$)^{6,9,10,16,17} or sufficient granulation tissue to allow skin grafting ($n=2$).⁵ The number of applications of citric acid ranged from 8 to 50, mainly on a daily basis but in some were on alternate days (Levels 4c & 4d).



There are film dressings ...

Then there's OPSITE[®] Post-Op



Smith & Nephew Pty Ltd
Australia
T 61 2 9857 3999 F 61 2 9857 3900
www.smith-nephew.com/australia
Customer Service
T 13 13 60 F 1800 671 000

[®]Trademark of Smith & Nephew
SNI2353 (02/16)

References

1. Tompkins L. OPSITE[®] Post-Op Visible Dressings Physical Properties. Data on file; 2010: report DS/10/084/R1. 2. Smith & Nephew Wound Management Laboratory Report. Data on file; 2009: report DS/08/149/R3. 3. Keene A. OPSITE Post-Op Visible (High MVTR) waterproof transparent dressing low allergy claim. Data on file. 4. Groves A, et al. An evaluation of a lightweight non-adherent dry dressing in the clinical situation. Data on file; 1983: UK 34.

Adverse effects

Application of citric acid to wounds can be associated with mild to moderate irritation for up to five minutes post application.^{3,4,8,10} (Levels 4b, 4c & 4d) In superficial burns the effect can be more severe in the first 2–3 minutes after application.¹ (Level 2b)

CHARACTERISTICS OF THE EVIDENCE

This evidence summary is based on a structured search of the literature and selected evidence-based healthcare databases (including WHO and WHO AFRO) using the terms citric acid and wounds. The evidence in this summary comes from:

- One double-blinded RCT¹⁴ (Level 1c)
- One quasi-experimental study¹ (Level 2c)
- Six descriptive observational studies^{2-4,7,8,13} (Level 4b)
- Four case series^{5,6,10,12} (Level 4c)
- Three case studies^{9,16,17} (Level 4d)
- Two laboratory studies^{11,15} (Level 5c)

BEST PRACTICE RECOMMENDATIONS

- Citric acid as a 3% irrigation solution or ointment applied daily offers a low cost option for managing chronic clinically infected wounds in settings where there is limited or no access to antibiotic therapy or where conventional treatment has been ineffective. (Grade B).
- Citric acid 3% could also be considered for treatment of limb wounds caused by trauma with delayed presentation for care. (Grade B)
- Assess patient's level of pain and provide pain relief if required prior to commencing dressing. (Grade A)

REFERENCES

1. Nagoba B, Gandhi R, Hartalkar A, Wadher B, Selkar S. Simple, effective and affordable approach for the treatment of burns infections. *Burns* 2010;36:1242–7.
2. Nagoba B, Gandhi R, Wadher B, Rao A, Hartalkar A, Selkar S. A simple and effective approach for the treatment of diabetic foot ulcers with different Wagner grades. *Int Wound J* 2010;7(3):153–8.
3. Nagoba B, Wadher B, Rao A, Kore G, Gomashe A, Ingle A. A simple and effective approach for the treatment of chronic wound infections caused by multiple antibiotic-resistant *Escherichia coli*. *J Hosp Infect* 2008;69:177–80.
4. Nagoba B, Gandhi R, Wadher B, Rao A, Selkar S. Simple and effective approach for the treatment of traumatic wounds in non-diabetic patients: a prospective open study. *Int Wound J* 2013;10:585–9.
5. Nagoba B, Gandhi R, Wadher B, Gandhi S, Selkar S. Citric acid treatment of necrotizing fasciitis: a report of two cases. *Int Wound J* 2010;7(6):536–8.
6. Nagoba B, Gandhi R, Suryawanshi N, Wadher B, Selkar S. Citric acid treatment of chronic non-healing ulcers in surgically excised cases of lateral malleolus bursitis. *Wound Prac Res* 2014;22(3):136–8.
7. Nagoba B, Wadher B, Rao A, Selkar S, Gandhi R. Treatment of lepromatous ulcers using citric acid as a sole antimicrobial agent. *Int Wound J* 2012;9:553–6.
8. Nagoba B, Wadher B, Gandhi R, Rao A, Selkar S, Hartalkar A. Citric acid treatment of surgical site infections: a prospective open study. *Wound Prac Res* 2011;19(2):82–6.
9. Nagoba B, Punpale A, Ayachit R, Gandhi R, Wadher B. Citric acid treatment of postoperative wound in an operated case of synovial sarcoma of the knee. *Int Wound J* 2011;8(4):425–7.
10. Nagoba B, Dawale C, Raju R, Wadher B, Chidrawar S, Selkar S *et al*. Citric acid treatment of post operative wound infections in HIV/AIDS patients. *J Tissue Viability* 2014;23:24–8.
11. Yabanoglu H, Basaran O, Aydogan C, Azap O, Karakayali F, Moray G. Assessment of the effectiveness of silver-coated dressing, chlorhexidine acetate (0.05%), citric acid (3%), and silver sulfadiazine (1%) for tropical antibacterial effects against the multi-drug resistant *Pseudomonas aeruginosa* infecting full-skin thickness burn wounds on rats. *Int Surg* 2013;98:416–23.
12. Nagoba B, Gandhi R, Wadher B, Potekar R, Kohle S. Microbiological, histopathological and clinical changes in chronic infected wounds after citric acid treatment. *J Med Microbiol* 2008;57(5):682–5.
13. Malu R, Nagoba B, Jaju C, Suryawanshi N, Maii S, Goyal V *et al*. Topical use of citric acid for wound bed preparation. *Int Wound J* 2014;1–4.
14. Prabhu V, Prasadi S, Pawar V, Shivani A, Gore A. Does wound pH modulation with 3% citric acid solution dressing help in wound healing: a pilot study. *Saudi Surg J* 2014;2(2):38–46.
15. Xie Z, Aphale N, Kadapure T, Wadajkar A, Orr S, Gyawali D *et al*. Design of antimicrobial peptides conjugated biodegradable citric acid derived hydrogels for wound healing. *J Biomed Mater Res* 2015;103A(12):3907–18.
16. Nagoba B, Punpale A, Podder A, Namdev M, Suryawanshi N, Swami G *et al*. Citric acid treatment of chronic nonhealing ulcerated tophaceous gout with bursitis. *Int J Lower Extremity Wounds* 2013;12(4):276–8.
17. Hartalkar A, Nagoba B, Wadher B. Citric acid treatment of large non-healing ulcer in a patient with chronic liver disease. *Eur J Gen Med* 2012;9(3):205–7.

Australian College of Nursing

26–28 October 2016

THE NATIONAL NURSING FORUM

THE POWER OF NOW

Melbourne Park Function Centre

ACN's signature event, provides a platform for nurses to learn from inspiring and preeminent nurse leaders, share insights and work together to advance nurse leadership, the nursing profession and health care outcomes for all Australians.

An extensive agenda covers the Power of the Nursing Workforce, the Power of Nursing Leadership, the Power of Nursing Care and the Power of Self Care.

Exploring and discussing four key streams, the National Nursing Forum looks at the current and most vital issues nurses face in clinical, management, education and leadership arenas and the ways in which nurses can lead the necessary reform to create a fitter and healthier health care system and nursing workforce.

For more information and to register
www.acn.edu.au/NNF2016 or contact events@acn.edu.au