

AUSTRALIAN + NEW ZEALAND
CONTINENCE JOURNAL

The management of urinary incontinence in nursing homes: a scoping review

Joan Ostaszkievicz, Leona Kosowicz, Jess Cecil, Deidree Somanader, Briony Dow

Improving ward-based neurogenic bladder and bowel care after acute spinal cord injury: a narrative review

Cristina Levy, Shilpa Gaikwad, Michael Whishaw, Wendy Bower

Abstracts

UroGynaecological Society of Australasia
Annual Scientific Meeting
Melbourne 22-25 November 2023



continence **NZ**

Contents

Editorial	79
The management of urinary incontinence in nursing homes: a scoping review	80
Improving ward-based neurogenic bladder and bowel care after acute spinal cord injury: a narrative review	101
Abstracts UroGynaecological Society of Australasia Annual Scientific Meeting Melbourne 22-25 November 2023	109
Australian news	112
New Zealand news	114



CONTINENCE FOUNDATION OF AUSTRALIA

Suite 1, 407 Canterbury Road,
Surrey Hills, VIC 3127
Tel +61 3 8692 8400
Fax +61 3 9347 2533
Email info@continence.org.au
Web www.continence.org.au

BOARD OF DIRECTORS

Chair — Elizabeth Compton
Lesley Barton
Celia Bolton
Prof Rosemary Calder
Dr Janet Chase
Jacinta Crickmore
Claire Richards
Dr Kathy Sloots
Company Secretary — Michael Corry
CEO — Rowan Cockerell



NEW ZEALAND CONTINENCE ASSOCIATION INC

PO Box 254, Waiuku 2341
New Zealand
Association Secretary
Tel +64 9238 3172
Freephone 0800 650 659
Email info@continence.org.nz
Web www.continence.org.nz

NATIONAL EXECUTIVE

President — Dr Anna Lawrence
Treasurer — Dr John Short (Acting)
Members — Liz Childs, Helen Peek,
Jacinta Townsend
CEO — Laura Fear

Summer 2023
Volume 29 Number 4
ISSN 2208-5750

THE OFFICIAL JOURNAL OF THE
CONTINENCE FOUNDATION OF
AUSTRALIA + THE NEW ZEALAND
CONTINENCE ASSOCIATION

Published four times a year by



10 Walters Drive
Osborne Park, WA 6017
www.cambridgemedia.com.au

Publisher Greg Paull
Copy editor Ceridwen Clocherty
Design and layout Gordon McDade

Advertising enquiries to
Simon Henriques
Cambridge Media
Tel (08) 6154 3912
Email simonh@cambridgemedia.com.au

© Continence Foundation of Australia
and New Zealand Continence
Association.

Apart from any use permitted under
the *Copyright Act 1968*, material
published in the *Australian and New
Zealand Continence Journal* may not be
reproduced without written permission
of the publisher.

The Journal is indexed with CINAHL,
Ebsco, infoRMIT.

ADVERTISING

Advertising that appears in the
*Australian and New Zealand
Continence Journal* conforms to the
standards required by the Continence
Foundation of Australia Ltd and the
New Zealand Continence Association,
but endorsement is in no way implied
by the publishing of said material. All
advertising enquiries should be directed
to the publisher, Cambridge Media.

EDITORIAL NOTE

The views expressed in the *Australian
and New Zealand Continence Journal* are
those of the authors and not necessarily
those of the Continence Foundation
of Australia Ltd, the New Zealand
Continence Association, the Editor,
the Editorial Committee or Cambridge
Media and must not be quoted as such.
While every care is taken to reproduce
articles as accurately as possible, the
publisher accepts no responsibility for
errors, omissions or inaccuracies.

Editorial Committee

Chair & Editor

Assoc Prof Christian Moro
BSc, BEd, MBus, PhD, SFHEA
Centre for Urology Research
Faculty of Health Sciences and Medicine, Bond University,
Gold Coast, QLD, Australia

Continence Foundation of Australia Representative

Mrs Rowan Cockerell, CEO

Continence New Zealand Representative

Laura Fear, CEO

Committee

Assoc Prof Wendy Bower
BAppSc (Physio), PhD, PgD Epidemiology and
Biostatistics
Physiotherapy Research Lead & Senior Continence
Clinician, The Royal Melbourne Hospital, Melbourne,
Victoria
Faculty of Medicine, Dentistry and Health Sciences,
The University of Melbourne, Melbourne, Victoria

Dr Anna Lawrence
MChB, FRACS (Uro)
Urologist, Genitourinary Reconstruction,
Auckland Spinal and Rehabilitation Unit, Auckland
Hospital, New Zealand

Prof Jean Hay-Smith
MSc PhD
Academic Head of the Rehabilitation Teaching and
Research Unit
Department of Medicine
University of Otago, Wellington

Ms Elizabeth Watt
RN, MNS, BAppSc (Adv Nur), Grad Cert Cont Prom, FACN
Senior Research Fellow (casual)
National Ageing Research Institute Ltd
Parkville, Melbourne, VIC, Australia

Prof Russ Chess-Williams
BSc(Hons), PhD, FBPhS
Director, Centre for Urology Research
Faculty of Health Sciences & Medicine, Bond University,
Gold Coast, QLD, Australia

Assoc Prof Vivien Wong
MBChB, FRANZCOG, CU, PhD
Urogynaecologist, Robina Hospital, Gold Coast, Robina,
QLD, Australia

Co-opted Members

Charlotte Phelps
BBIomedSci(Hons), PhD Candidate in Urology
Centre for Urology Research
Faculty of Health Sciences and Medicine, Bond University,
Gold Coast, QLD, Australia

Prof Kate Moore AO
MBBS, MD, FRCOG, FRANZCOG, CU
Urogynaecologist, St George Hospital, Kogarah,
NSW, Australia

Dr Vincent Tse
MB, BS (Hons), MS (Syd), FRACS
Urologist, Concord Hospital, Sydney, NSW, Australia

Managing Editor

Sarah Taylor
Continence Foundation Australia
Suite 1, 407 Canterbury Road, Surrey Hills,
VIC 3127, Australia
Tel 03 8692 8400
Email journal@continence.org.au

Call for Papers

The *Australian and New Zealand Continence Journal* seeks articles and original research papers from people practising and researching the management and treatment of incontinence and continence health promotion.

Do you need topic ideas? A variety of topics are possible and include, but are not limited to: outcome studies, aged care, paediatrics, pregnancy and childbirth, novel drug therapies, reviews of devices, either surgical or non-surgical, assessment articles, literature reviews of continence-related topics, home and community care issues and successes, men's health, nursing management, physiotherapy management, support by other allied health disciplines (including occupational therapy and social workers), the psychological impact of living with incontinence, ethical issues, cultural issues and collaborative approaches to care.

Articles may be papers for peer review, clinical updates, case studies or evaluation of programs.

To discuss topics or for assistance in the preparation of papers and articles, please email journal@continence.org.au

EDITORIAL

For referencing Moro C. Editorial. Australian and New Zealand Continence Journal 2023; 29(4):79.

DOI <https://doi.org/10.33235/anzcj.29.4.79>

Licensed under CC BY 4.0

It is with great pleasure that we present the final edition of the *Australian and New Zealand Continence Journal for 2023*. Our first article, *The management of urinary incontinence in nursing homes: a scoping review*¹, identifies the various interventions employed for the management of urinary incontinence in nursing homes. This article not only provides an excellent summary of the various policies and practices, but makes recommendations regarding which approaches ensure the most benefit. We are confident that this will be of great interest to many of our readers. The second article in this edition, *Improving ward-based neurogenic bladder and bowel care after acute spinal cord injury: a narrative review*², focuses on the procedures, protocols and processes that occur after spinal cord injury. After investigating the literature, this article provides recommendations for management and insights into processes that can particularly assist in the acute phase of spinal cord injury.


In this edition we have been very happy to collaborate with the UroGynaecological Society of Australasia to publish a small selection of abstracts from the Annual Scientific Meeting (22-25 November 2023, Melbourne). This includes: Clinical Quality Indicators using Australasian Pelvic Floor Procedure Registry clinical data³; Biomechanical and immunobiological properties of human fascia lata (HFL) vs TiMesh®: Implications for pelvic reconstructive surgery⁴; and Mesh perforation into viscus following pelvic mesh surgery: experiences and recommendations for diagnosis and management⁵. It is absolutely fantastic to see so much great research being conducted that relates to the pelvic region, and overall continence.

I express a warm welcome to Sarah Tayler who has taken on the managing editor role for the journal. Sarah has a background in editing, writing, marketing and production, and it has been absolutely amazing to have her on board as part of our journal's team. Sarah's experiences will greatly assist as we strategically develop and grow the journal throughout 2024. This includes working to increase the visibility of published articles, promoting the research presented within, and enhancing our engagement with the broader research community. We very much appreciate the support received from our authors, reviewers and members of the editorial committee. If you would like to express interest in reviewing or need a hand preparing or submitting an article, please contact us at journal@continence.org.au

The ANZCJ presents an excellent avenue for the publication of research. All articles are available online as diamond open access, with no cost to authors or readers. Each publication is allocated an individual digital object identifier (DOI) to assist with referencing and tracking, and is listed across several databases. If you are a first-time author, our reviewing and editorial teams can offer guidance, feedback and support to assist. Performing and publishing research not only helps our community with increased knowledge and insights, but also advances the vision of the Continence Foundation of Australia and Continence New Zealand to promote bladder and bowel health and eliminate the stigma and restrictions of all aspects of incontinence.

REFERENCES

1. Ostaszkiwicz J et al. The management of urinary incontinence in nursing homes: a scoping review. Australian and New Zealand Continence Journal 2023; 29(4):80-100.
2. Levy C et al. Improving ward-based neurogenic bladder and bowel care after acute spinal cord injury: a narrative review. Australian and New Zealand Continence Journal 2023; 29(4):101-108.
3. Kartik A. Clinical Quality Indicators using Australasian Pelvic Floor Procedure Registry clinical data. Australian and New Zealand Continence Journal 2023; 29(4):109.
4. Hennes D. Biomechanical and immunobiological properties of human fascia lata (HFL) vs TiMesh®: implications for pelvic reconstructive surgery. Australian and New Zealand Continence Journal 2023; 29(4):110.
5. Tan YH. Mesh perforation into viscus following pelvic mesh surgery: experiences and recommendations for diagnosis and management. Australian and New Zealand Continence Journal 2023; 29(4):111.



Christian Moro
Editor-In-Chief and Chair
Australian and New Zealand
Continence Journal

The management of urinary incontinence in nursing homes: a scoping review

For referencing Ostaszkiwicz J et al. The management of urinary incontinence in nursing homes: a scoping review. Australian and New Zealand Continence Journal 2023; 29(4):80-100.

DOI <https://doi.org/10.33235/anzcj.29.4.80-100>

Licensed under CC BY 4.0

Submitted 3 October 2023, Accepted 4 November 2023

ABSTRACT

The aim of the research was to identify interventions for the management of urinary incontinence (UI) in nursing homes. A scoping review was conducted with methods adapted from *The Joanna Briggs Institute reviewers' manual 2015 methodology for conducting scoping reviews* and reporting was guided by the *Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for scoping reviews guidelines*. Findings were synthesised without meta-analysis. Databases that were searched were MEDLINE, Embase, PsychINFO, CINAHL, JBI Evidence-Based Practice Database and the Cochrane Database of Systematic Reviews from 2010 to September 2021. The search was augmented with hand searching.

A total of 3,885 records were located. After exclusions and screening of 370 full-text articles, 30 publications were included – seven systematic reviews, 15 randomised-controlled trials, seven quasi-experimental studies and one cohort study. Studies addressed toileting assistance programs, exercise programs, drug therapies, technology-based interventions, education programs and multi-component interventions.

Multi-component interventions facilitated by specialist healthcare professionals offer the strongest evidence. Evidence about exercise programs was limited and inconsistent, as was evidence for anticholinergics. Transcutaneous posterior tibial nerve stimulation is unlikely to reduce rates of UI in nursing homes with residents with high rates of cognitive impairment. Further evidence is required on the use of telemonitoring systems. Education programs for staff that provide on-site support and competency-based learning, led by specialist healthcare professionals, improve staff knowledge, attitudes and compliance with assessments, toileting and documentation. Education about person-centred approaches is required to provide appropriate care for residents living with dementia.

We conclude that a multicomponent approach led by a specialist such as a nurse with advanced

clinical and leadership skills offers the most benefit. Nursing home policies and practices should focus on education programs for staff, with interventions that increase residents' choice, activity level, nutritional status, hydration and toileting opportunities.

Keywords scoping review, urinary incontinence, nursing homes, residential aged care, evidence

INTRODUCTION

Urinary incontinence (UI) in nursing homes has physical, psychological, economic and social consequences for residents, their families and staff. Despite its impact, little is known about strategies for the organisation and delivery of care. Medically, incontinence is conceptualised as a symptom of an underlying condition that can be prevented and treated, even in frail older people in nursing homes¹. International research suggests 50–70% of nursing home residents have UI^{2–5},

Joan Ostaszkiwicz*

National Ageing Research Institute, VIC, Australia
Federation University, Health and Innovation Transformation Centre, VIC, Australia
University of Melbourne, Faculty of Medicine, Dentistry and Health Sciences, VIC, Australia

Leona Kosowicz

National Ageing Research Institute, VIC, Australia

Jess Cecil

National Ageing Research Institute, VIC, Australia

Deidree Somanader

Monash University, Faculty of Medicine, Nursing and Health Sciences, VIC, Australia

Briony Dow

National Ageing Research Institute, VIC, Australia
University of Melbourne, Faculty of Medicine, Dentistry and Health Sciences, VIC, Australia

*Corresponding author

Conflict of interest

The authors declare no conflicts of interest.

Funding

The scoping review was funded by the Continence Foundation of Australia as part of a research study to develop a best practice model of continence care for residential aged care homes in Australia.

making it an important policy and practice issue for nurses and the nursing profession. In many countries, the quality of continence care in institutional settings is not consistent with recommended prevention and treatment guidelines and falls far below the public's expectations, as evidenced by the Mid Staffordshire NHS Foundation Trust Public Inquiry⁶ in the UK and the Royal Commission into Aged Care Quality and Safety⁷ in Australia. However, understandings about what constitutes quality continence care in nursing homes vary^{8,9}. As such, an investigation summarising the quantitative evidence about interventions for UI management is required.

THE REVIEW

The causes of UI in nursing homes are multifactorial¹. The International Consultation on Incontinence identifies both intrinsic (individual) and extrinsic (environmental) factors that increase nursing home residents' risk of developing UI¹. There is a mounting body of evidence to suggest that the processes of care in nursing home settings are not designed to promote therapeutic continence care for care-dependent older people⁶⁻¹⁰.

A modelling of risk factors for UI in nursing home residents living with dementia found the strongest predictor was impaired functional status, expressed and measured as dependence in activities of daily living (ADL)¹¹. This is an important finding that highlights the need for interventions to optimise nursing home residents' functional status and prevent premature decline.

Another potential cause of UI in nursing homes is a lack of assessment to identify potentially reversible causes. An early study from the US found up to 81% of nursing home residents had potentially reversible causes of UI that were rarely investigated or treated¹².

UI has a negative impact on some domains of nursing homes residents' quality of life (QOL)¹³. Xu and Kane¹³ compared the overall QOL of 8,620 eligible residents with and without UI from 371 nursing homes. Although UI was not associated with overall QOL, it decreased the QOL domains of dignity, autonomy and mood. Jerez-Roig et al.¹⁴ found 24.1% of residents with UI reported a severe impact on their QOL.

UI also impacts on nursing home residents' physical health, increasing their risk of incontinence-associated dermatitis¹⁵, falls¹⁶, functional decline¹⁴ and death¹⁷. Organisational costs are another important consideration. Research from the US from 2003 reported that the cost of proactively managing UI in nursing homes over a 6-month period was US\$586,436, with 46% attributed to direct labour costs alone¹⁸.

Toileting assistance programs have been the focus of prior research on the management of UI in nursing homes. Seminal research conducted by Schnelle¹⁹⁻²³ demonstrated it was possible to reduce rates of UI immediately following the implementation of a toileting assistance program in this setting. The

challenge has been for usual care staff to sustain these improvements in usual care conditions²⁴. There is a lack of contemporary data about actual rates of toileting assistance programs in nursing homes, as well as evidence about the best ways to identify residents who would benefit from this intervention. Similarly, there are questions about the best ways to implement and sustain toileting assistance programs for residents who need this assistance.

To address the personal, social and financial costs of UI in nursing homes there is a need to better understand the range of interventions available and their evidence base. This information could inform workforce planning, research, future funding and policies and practice for nursing home care.

Aim

The aim of this scoping review was to map and summarise the body of quantitative evidence about interventions for the management of UI in nursing homes. The PICO question was What interventions have been trialled for the management of UI in nursing homes and what is their effectiveness?

METHODS

Design

A scoping review was conducted using methodology adapted from *The Joanna Briggs Institute (JBI) reviewers' manual 2015 methodology for JBI scoping reviews*²⁵ and reporting methods were guided by *PRISMA-ScR (Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for scoping reviews) guidelines*²⁶. The rationale for conducting a scoping review rather than a systematic review was to map research about interventions for the management of UI in nursing homes, ie, to identify interventions that have been evaluated using quantitative research methods.

Search methods

The bibliographic search of health and medical databases followed a three-step methodology described by the JBI guidelines. Firstly, limited searches of Ovid MEDLINE and CINAHL were conducted. An experienced librarian was consulted to assist with identifying relevant keywords from the titles and abstracts of the articles retrieved from this initial search. Secondly, a comprehensive search for English-language articles published from 2010 to September 2021 was performed using the finalised keywords and MeSH terms. Six electronic databases were comprehensively searched – MEDLINE, Embase, PsychINFO, CINAHL, JBI Evidence-Based Practice Database, and the Cochrane Database of Systematic Reviews. The search strategy is available in Table 1. Lastly, selected publications were hand searched for additional relevant references.

Inclusion and exclusion criteria

The search was limited to primary sources and systematic reviews of primary sources. Grey literature

and government reporting was excluded. Publications that evaluated interventions for the management of UI in nursing home residents were included. Publications that focused on the management of other types of incontinence (ie, faecal or dual incontinence), incontinence-associated dermatitis, catheter-associated urinary tract infections, urinary catheters or ostomies were excluded. Surgical interventions for the management of UI were also excluded.

Search outcome

All identified articles were imported into Covidence for screening (www.covidence.org). Three reviewers

(LK, JO and DS) independently reviewed the titles and abstracts of all identified sources from the database search. Five reviewers (LK, JO, BD, JC and DS) then completed a full-text review of the remaining material, with conflicts resolved through consensus.

Data abstraction

A data extraction form was developed to identify the key characteristics of each study as well as relevant information regarding interventions for the management of UI in nursing homes. Three reviewers (LK, JO and DS) independently extracted the data and resolved inconsistencies through consensus. The variables included study objectives, methods, setting/sample characteristics, intervention type, outcomes measured, findings, and key recommendations.

Synthesis

Because of anticipated heterogeneity of included studies, findings were synthesised without meta-analysis²⁷. They were then reported in narrative form according to the type of intervention, ie, toileting assistance programs, exercise programs, drug therapies, technology-based interventions, education programs, or multicomponent interventions.

RESULTS

Search outcomes

A total of 3,885 records were located through the database search, and one additional study was found through citations. Of the publications identified, 30 were included in the final analysis (Figure 1). This included seven systematic reviews²⁸⁻³⁴, 15 randomised-controlled trials (RCTs)³⁵⁻⁴⁹, seven quasi-experimental studies⁵⁰⁻⁵⁶ and one retrospective cohort study⁵⁷. The majority of intervention studies were from Europe (n=10, 43.6%), while the remaining studies were from the US (n=7, 30.5%), Asia (n=3, 13%), Australia (n=1, 4.3%), New Zealand (n=1, 4.3%) and Iran (n=1, 4.3%).

Characteristics of included studies

Of the 30 included publications, six evaluated toileting assistance programs^{29,31,32,42,46,55}, four addressed exercise programs^{29,34,47,48}, three addressed drug therapies^{29,33,57}, and two described a technology-based intervention^{37,56}. A total of 11 publications described the effects of education programs for nursing home staff^{38-41,44,45,49,51-54} and six evaluated multicomponent interventions^{29,28,35,36,43,50}. There was considerable heterogeneity with respect to the types of interventions, how they were delivered (ie, duration, intensity, components), and how outcomes were evaluated. Therefore, there is no one intervention that can be recommended over another. The following section describes the evidence for each intervention type; detailed characteristics of the included studies are presented in Tables 2-5.

Toileting assistance programs

Systematic reviews from 2011³¹, 2012²⁹ and 2015³² reported moderate evidence that toileting assistance

Table 1. Search strategy

#	Query	Limiters/ expanders
S1	aged care or nursing home or residential aged care facility or long term care	Limiters: English language Published date: 20100101-20201231 Expanders: apply equivalent subjects Search modes: Boolean/Phrase
S2	(continence or incontinence) OR continence management OR continence assessment OR continence promotion OR continence care OR continence support OR continence assessment tools OR (incontinence or incontinent or urinary incontinence or urinary incontinent or urinary leakage) OR incontinence care OR incontinence management OR bladder management OR toileting program OR toileting assistance OR toileting intervention	
S3	care dependent OR care dependency	
S4	dementia or Alzheimer's or cognitive impairment	
S5	Residents' perspectives OR residents' experiences OR residents' views OR residents' opinions OR residents' attitudes	
S6	(staff perceptions or attitudes or opinions or experiences or views or reflection or beliefs) OR staff perspectives	
S7	(family experience or family perspectives or relatives experiences) OR families perceptions OR family opinions OR family attitudes	
S8	S1 AND S2	
S9	S8 AND S3	
S10	S8 AND S4	
S11	S8 AND S5	
S12	S8 AND S6	

programs can reduce the frequency of UI in nursing homes, increase residents' appropriate toileting behaviour, maintain their social continence, and regain continence. However, they also reported that the effects were short-term and dependent on staff resources and time³².

A further three studies suggest it is possible to implement toileting assistance programs in nursing homes over a longer period of time and with usual care staff. These include a RCT conducted by Lai and Wan⁴² that evaluated a 6-month, 2-hourly prompted voiding (PV) intervention implemented by usual care staff in 52 nursing home residents in five nursing homes in Hong Kong. The intervention was compared with usual care. The researchers reported a significant difference in wet episodes per day ($p=0.001$), UI rate per day ($p=0.001$), and total continent toileting per day ($p\leq 0.001$) with positive results in the intervention group. Using a quasi-experimental design, Suzuki et al.⁵⁵ trialled PV for 1 week followed by ultrasound-assisted prompted voiding (USAPV) for 12 weeks in a cohort of 77 nursing homes residents in six nursing homes in Japan. The researchers reported an 11.8% reduction in the cost of incontinence products from baseline ($p=0.006$) and significant improvements in care workers' QOL.

In 2019, the same authors conducted an RCT comparing PV alone with USAPV in 13 nursing homes with 125 residents over an 8-week period⁴⁶. They

reported a significant reduction in daytime UI for the USAPV group than for the PV group ($p=0.018$), with the proportion of residents whose daytime urine loss decreased by greater than 25% being 51% and 26% in the USAPV and PV group respectively ($p=0.020$). Furthermore, while carer burden was unchanged in the USAPV group, it significantly worsened in the PV group after the intervention ($p=0.010$).

Technology-based interventions

One randomised placebo-controlled trial evaluated the effectiveness of a technology-based intervention, namely transcutaneous posterior tibial nerve stimulation³⁷. A total of 12 30-minute sessions of this nerve stimulation were provided to 197 nursing home residents with UI over 6 weeks, while 209 residents were provided with sham stimulation. The researchers reported a statistically significant greater reduction in UI over 24 hours in the sham group in comparison to the treatment group ($p=0.05$). Therefore, this study did not demonstrate evidence that transcutaneous posterior tibial nerve stimulation is an effective intervention for UI.

A quasi-experimental study by Yu et al.⁵⁶ evaluated the effectiveness of a telemonitoring system. In this study, a sensor device was placed in residents' incontinence products to detect the onset of wetness episodes in 31 residents over a 72-hour assessment period. The data




Figure 1. PRISMA diagram of study selection process

obtained from the assessment was used to develop an individualised continence care plan for each resident. The researchers reported a significant reduction in the volume of urine voided into the pads ($p=0.015$), significant increases in the number of prescribed toileting visits ($p=0.015$), significant increases in the actual number of toilet visits ($p\leq 0.001$), successful toileting events ($p=0.011$), and in staff adherence to urinary continence care plans ($p=0.033$).

Exercise programs

In an RCT, Tak et al.⁴⁷ evaluated the effects of a 6-month group-based behavioural exercise program on 102 female nursing home residents from 20 nursing homes in the Netherlands. Although physical performance significantly improved for the intervention group compared to the control group ($p\leq 0.001$), no significant differences were found for continence-related or QOL outcomes. In contrast, when personalised treatment goals were designed for each resident and physical activity and ADL training was implemented for 3 months, Vinsnes et al.⁴⁸ reported a significant decrease in the amount of UI leakage among 35 nursing home residents in Norway ($p=0.03$). One systematic review reports exercise and functional incidental training programs reduced wetness rates and improved rates of appropriate toileting²⁹. However, in a subsequent review, Stenzelius et al.³⁴ reported a very low quality of evidence for physical activity in combination with ADL training in reducing UI.

Drug therapies

Two systematic reviews were identified that evaluated the use of drug therapies for the management of UI in nursing homes. In one review, anticholinergics showed a small but significant positive effect on urinary leakage in urge UI³³. However, the review also found that adverse events, including dry mouth and constipation, were common, and studies on the impact of anticholinergics on cognition in this population have not yet been conducted. It was concluded that the use of anticholinergics for the treatment of urge UI in frail older people is not evidence-based. Flanagan et al.²⁹ reported that the use of anticholinergics (oxybutynin) as an adjunct to toileting therapy may have benefits for the reduction of UI, but benefits appear to be inconsistent and may be small or non-existent.

One cohort study also evaluated the use of drug therapies for the management of UI, namely bladder antimuscarinics (BAMs)⁵⁷. Through new user design, BAM users were compared with propensity score (PS)-matched non-users, and the benefits (change in UI status and QOL) and harmful effects (fracture risk and decline in cognitive performance) of BAM initiation were evaluated. While BAM initiation was significantly associated with improvement in UI (OR=1.34, 95% CI 1.13-1.60), the risk of fractures was also significantly increased in new users compared to non-users (hip fracture: hazard ratio [HR]=3.67, 95% CI 1.46-9.34; 'any' fracture: HR=2.64, 95% CI 1.37-5.10) and there was no clinically significant improvement in social engagement among BAM users. It was concluded that

non-selective immediate release BAMs (especially immediate-release oxybutynin chloride) should be approached cautiously as a treatment for UI amongst the nursing home population.

Education programs

Of the 11 publications that evaluated the effects of education programs for nursing home staff, seven were RCTS^{38-41,44,45,49} and four were quasi-experimental studies⁵¹⁻⁵⁴. The nature of the educational interventions and the outcomes of interest varied considerably.

Over a 10-month period, Wijk et al.⁴⁹ implemented an education program about a person-centred approach to continence care for nursing home staff from two nursing homes in Switzerland. Staff in the intervention group were educated on how to elicit information from residents about their personal identity and preferences and how to use this information to design an individualised continence care plans for each resident. In homes where the approach was used, the number of times residents received toileting assistance was significantly higher during and after the intervention compared to the control home ($p=0.019$). A significant positive effect was also found in the number of continence assessments conducted in the intervention homes ($p\leq 0.05$).

Also conducted in seven Swiss nursing homes over a 14-month period was a step wedge study led by Kohler et al.⁴¹. The intervention was a 4-hour education program focusing on continence care for residents living with dementia followed by six case conferences (one per month). The researchers found a decrease in UI for all groups which was only significant in the 11-month group ($p=0.036$). QOL also significantly increased in seven of the nine domains for residents in all groups.

In Austria, Hödl et al.⁴⁰ conducted a cluster RCT in which managers from 12 nursing homes received and discussed 29 recommendations for the conservative management of UI. Three months later, residents in the intervention sites were found to have a significantly lower risk of daily UI ($p=0.02$), were less likely to receive absorbent products ($p=0.01$), and were five times more likely to receive the recommended interventions ($p\leq 0.001$) than those who received usual care.

Two further publications were identified, related to one large cluster RCT conducted across eight nursing homes in four European countries – Sweden, England, Ireland and the Netherlands^{39,45}. The aim was to compare standard dissemination of evidence-based guidelines about UI with two different types of facilitation. The guidelines promoted: (i) screening for UI; (ii) a detailed continence assessment; (iii) an individualised treatment plan; and (iv) referral if needed. The researchers reported no between-group significant differences in the documented percentage compliance with the continence recommendations, which was assessed at baseline, and at 6, 12, 18 and 24 months after the intervention and noted that all study arms improved over time. It was not possible

Table 2. Characteristics of included studies: systematic reviews

Legend for all tables:
 NH: nursing home; UI: urinary incontinence; FI: faecal incontinence; IG: intervention group; CG: control group
 NA: nursing assistant; CNA: certified nursing assistant; CNS: clinical nurse specialist; IF: internal facilitator; LPN: licensed practical nurse

Review objectives	Descriptions of interventions/ phenomena of interest	Methods	Findings
Donald et al. (2013)²⁸ To report quantitative evidence of the effectiveness of advanced practice nursing roles, clinical nurse specialists (CNS) and nurse practitioners (NP) in meeting the healthcare needs of older adults living in long-term NH settings	Evaluation of NP and/or CNS in long-term care Outcome measures: functional status, health status, QOL, health services use, and resident and family satisfaction	Quantitative systematic review	8,277 studies located, 15 papers about four studies included (all interventional studies) Long-term care settings with advanced practice nurses had lower rates of depression, UI, pressure ulcers, restraint use, and aggressive behaviours, more residents who experienced improvements in meeting personal goals, and family members who expressed more satisfaction with medical services Interventions that reduced or slowed decline in UI: <ul style="list-style-type: none"> CNS facilitation of evidence-based protocols and CNS provision of staff education, consultation, and direct care to residents for 6 months post-admission As above, plus CNS membership on the NH's quality assurance committee; CNS provision of formal in-services to certified nursing assistants (CNA); and CNS collaboration with staff in problem solving teams
Flanagan et al. (2010)²⁹ To describe and compare interventions for the management of incontinence and promotion of continence in care home residents	UI, bladder training, PV, timed voiding, habit training, functional incidental training, and prompted walking program	Systematic review of studies investigating urinary and faecal incontinence (FI) in people aged 65 years or older in care homes	181 studies located, 68 included (42 interventional studies, 26 observational/descriptive studies). 3,224 residents total, mainly female, aged >65 years Most common treatments: <ul style="list-style-type: none"> Toileting programs and incontinence pads (effective in reducing frequency of incontinence, increasing appropriate toileting behaviour, helping to maintain social continence, and helping to regain continence) Drug therapy adjunct to toileting programs (inconsistent benefits) Combined physical therapy/behavioural therapies (short-term improvements) Adaptations to physical environment and staff training Exercise and Functional Incidental Training programs (reduce wetness rates and improve appropriate toileting rates) Combined complex behavioural interventions (effectiveness not yet determined)
Ostaszkievicz et al. (2020)³⁰ To describe, critique, and summarise research about the effects of education about UI on nurses' and nursing assistants (NAs)' knowledge and attitudes toward UI, their continence care practices, and patient outcomes	UI educational intervention that reported pre- and post-quantitative data about the effects of education programs designed to evaluate or improve four outcomes: <ul style="list-style-type: none"> Nurses' and/or NAs' knowledge of UI Nurses' and/or NAs' attitudes about UI Nurses' and/or NAs' continence care practices Patient outcomes 	A systematic review of randomised controlled studies or non-randomised or quasi-randomised trials	Conducted in NH in the US (19 trials, enrolled 1301 participants: 911 nurses, 235 NAs and 155 unclear) Findings: <ul style="list-style-type: none"> Nurses and NAs had limited ability to determine UI type and factors that require assessment, interpret clinical data to make a diagnosis, and identify patients who may be suitable for active interventions Increased knowledge about UI does not necessarily translate into improvements in practices that improve patient outcome A key barrier to the uptake of educational recommendations about UI could be a lack of audit and feedback data about patients' objective continence status

Review objectives	Descriptions of interventions/ phenomena of interest	Methods	Findings
<p>Roe et al. (2011)³¹</p> <p>To undertake a narrative review of descriptive empirical studies on the management of incontinence, promotion of continence or maintenance of continence in people aged 65 years and above in care homes with UI as the primary focus</p>	<p>Descriptive, observational or intervention studies (which include nursing, medical or behavioural interventions) aimed at the management of incontinence or the promotion or maintenance of continence; surgical or pharmacological interventions excluded</p>	<p>A systematic review of studies that used quantitative or qualitative designs and methods was undertaken to provide a narrative synthesis</p>	<p>167 located, 60 included (37 intervention studies, 23 descriptive observation studies). Participants were mainly women, mean age >80 years</p> <p>Findings:</p> <ul style="list-style-type: none"> • Prevalence of UI was higher than FI with more women affected than men • Use of incontinence pads and toileting programmes comprised the most common management approaches used • The evidence base, theories underpinning toileting practices or their form, frequency and content are unclear from the studies reviewed but are stated to comprise bladder training, scheduled or PV
<p>Roe et al. (2015)³²</p> <p>To synthesise evidence from systematic reviews on the management of UI and promotion of continence using conservative/behavioural approaches in older people in care homes to inform clinical practice, guidelines and research</p>	<p>Conservative approaches for continence promotion, continence maintenance or management of incontinence (excluding drugs or surgical interventions) from behavioural or nursing perspectives</p>	<p>A systematic review of systematic reviews with narrative synthesis</p>	<p>42 located, five included (all systematic reviews), 535,178 residents total mainly females, mean age range 73.9–88.7 years</p> <p>Findings:</p> <ul style="list-style-type: none"> • Toileting programmes, PV, use of incontinence pads are the main conservative behavioural approaches (evidence of effectiveness in the short-term dependent on staff adherence and resources) • Evidence from associated factors (exercise, mobility, comorbidities, hydration, skin care, staff perspectives, policies, and experiences and preference) are limited
<p>Samuelsson et al. (2015)³³</p> <p>To systematically review the efficacy of pharmacological treatment for UI in the elderly and frail elderly</p>	<p>Pharmacological treatment for UI in the frail elderly population</p>	<p>A systematic review of prospective controlled trials of pharmacological treatment for UI in persons aged 65 years or over</p>	<p>1045 located, 13 included (all placebo-controlled randomised studies), Frail elderly (three studies) and elderly (10 studies)</p> <p>Findings:</p> <ul style="list-style-type: none"> • Anticholinergic drugs have a small, but significant, effect on urinary leakage in the elderly with urgency UI • Adverse effects (dry mouth, constipation) were common. No studies assessed effect on cognition • Treatment with anticholinergics for urgency UI in the frail elderly is not evidence-based
<p>Stenzelius et al. (2015)³⁴</p> <p>To investigate the evidence and the effect of conservative treatment of UI and the QOL among older and frail older persons</p>	<p>Conservative treatment including complex interventions, toilet training, behaviour therapy, PV, environmental, complementary therapies, pelvic floor exercise, bladder training and electrical stimulation</p>	<p>A systematic review of randomised controlled studies and prospective, non-randomised studies of interventions of conservative treatment of UI in an older population (65 years or older)</p>	<p>1711 located, nine included (eight studies, three meta-analyses), four studies in NHs</p> <p>Results for NHs:</p> <ul style="list-style-type: none"> • Attention training (i.e., being reminded to make toilet visits and being helped to go to toilet every hour) in combination with functional exercise reduced urinary leakage episodes (moderate quality of evidence) • Physical training (i.e., physical activity) in combination with ADL training had no significant effect on urinary leakage (very low quality of evidence)

Table 3. Characteristics of included studies: RCTs

Aim(s)	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Azizi et al. (2019)³⁵</p> <p>To examine the effect of a self-care programme on UI and self-esteem in elderly men dwelling in NHs in Iran</p>	<p>n=61 residents (100% male, mean age 68.18 years, UI score ≥3 [severe UI]) n=2 NHs in Iran</p>	<p>Eight 45–60 minute weekly sessions comprising:</p> <ul style="list-style-type: none"> • Light strengthening exercises • Pelvic floor exercise • Lectures, group discussions, roleplay and educational booklet about UI improvement and the role of behavioural and lifestyle factors in the health of the urinary system <p>IG: self-care programme for 8 weeks plus 4-week follow-up (n=30) CG: usual care for 12 weeks (n=31)</p>	<p>UI severity (ICIQ-SF) Self-esteem (Rosenberg Self-Esteem Scale)</p>	<ul style="list-style-type: none"> • Significant decrease in the mean scores of ICIQ-SF in the IG compared with CG following intervention (ps0.001) • Significant increase in self-esteem in IG compared to CG (ps0.001) • No reported drop out
<p>Boorsma et al. (2011)³⁶</p> <p>To determine the effects of multidisciplinary integrated care on the quality of care and QOL for elderly people in NHs</p>	<p>n=462 residents randomised (IG 76.1% female, mean age 85.8 years; CG 74.1% female, mean age 85.5 years; all with physical or cognitive disabilities) n=10 NHs in the Netherlands</p>	<p>Geriatric assessment of functional health every 3 months comprising:</p> <ul style="list-style-type: none"> • Use of the Long-term Care Facility version of the Resident Assessment Instrument by trained nurse assistants to guide the design of an individualised care plan • Discussion of outcomes and care priorities with the family physician, the resident and his or her family • Monthly multidisciplinary meetings with the nurse assistant, family physician, psychologist and geriatrician to discuss residents with complex needs <p>IG: multidisciplinary integrated care (n=201) CG: usual care (n=139)</p>	<p>Primary outcomes:</p> <ul style="list-style-type: none"> • Sum score of 32 risk-adjusted quality-of-care indicators • Health-related QOL (short-form Rand Health Insurance Study questionnaire) <p>Secondary outcomes:</p> <ul style="list-style-type: none"> • 32 individual risk-adjusted quality-of-care indicators • ADLs (Groningen Activity Restriction Scale) • Quality of care (QUOTE-Elderly instrument short form) • Hospital admissions recorded at the (single) local hospital • Mortality 	<p>Compared with the facilities that provided usual care, IG facilities had a significantly higher sum score of the 32 quality-of-care indicators (mean difference -6.7, p=0.009; a medium effect size of 0.72). They also had significantly higher scores for 11 of the 32 indicators of good care in the areas of communication, delirium, behaviour, continence, pain and use of antipsychotic agents</p>
<p>Booth et al. (2021)³⁷</p> <p>To determine the clinical effectiveness of transcutaneous posterior tibial nerve stimulation to treat UI in care home residents and to determine the associated costs of the treatment</p>	<p>n=406 residents randomised (mostly female, mean age 85 years; UI at least weekly) n=37 NHs in UK</p>	<p>12 30-minute sessions of transcutaneous posterior tibial nerve stimulation (TPTNS) IG: TPTNS for 6 weeks (n=197) Sham group: Sham stimulation for 6 weeks (n=209)</p>	<p>Primary outcome:</p> <ul style="list-style-type: none"> • Change in volume of urine leaked over a 24-hour period at 6 weeks <p>Secondary outcomes:</p> <ul style="list-style-type: none"> • Number of pads used • Perception of bladder condition • Toileting skills • QOL • Resource use 	<p>Primary outcome:</p> <ul style="list-style-type: none"> • Primary intention-to-treat adjusted analysis indicated a mean change of -5ml (standard deviation 362ml) urine leakage from baseline in the IG and -66ml (SD 394ml) urine leakage in the sham group, which was a statistically significant, but not clinically important, between-group difference of 68ml urine leakage (95% CI 0-136ml; p=0.05) in favour of the sham group <p>Secondary outcomes:</p> <ul style="list-style-type: none"> • No meaningful differences were detected

Aim(s)	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Gencbas et al. (2018)³⁸</p> <p>To assess the effectiveness of the nursing care plan implemented using NNN linkages for elderly women with incontinence and living in NHs</p>	<p>n=62 residents randomised (100% female, 43.8% IG and 53.3% of CG in age range 75–85 years, MMSE score >23)</p> <p>n=5 NHs in Turkey</p>	<p>Staff received education on the use of three Standard Nursing Languages (NANDA, NOC, NIC) and implemented these in incontinence care plans</p> <p>IG: residents received care from SNL trained staff for 12 weeks (n=32)</p> <p>CG: usual care for 12 weeks (n=30)</p>	<p>Pre- and post-intervention measures:</p> <ul style="list-style-type: none"> Five Nursing Outcomes Classification (NOC) scales (“0503 urinary elimination”, “0502 urinary continence”, “1101 tissue integrity: skin and mucous membranes”, “0310 self-care toileting” and “medication response”) Incontinence Severity Index (ISI) Urinary Distress Inventory-6 (UDI-6) Incontinence QOL (I-QOL) scale 	<ul style="list-style-type: none"> Statistically significant improvement in all NOC scales in IG compared to the CG (urinary elimination p=0.001; urinary continence p=0.001; tissue integrity p=0.001; self-care toileting p=0.001; medication response p=0.004) Significant and positive difference was found in terms of the ISI (p=0.001), UDI-6 (p=0.001), and I-QOL (p=0.001) scale scores for the IG group post-intervention
<p>Harvey et al. (2012)³⁹</p> <p>To describe the process of introducing PACES to internal facilitators (IFs)</p> <p>To discuss the progress made during a 12-month period of collecting and analysing audit data using PACES</p> <p>To summarise the collective experience of using PACES, including reflections on its strengths and limitations</p>	<p>n=8 NHs in Sweden, England, Ireland and the Netherlands</p>	<p>Intervention: a 3-day workshop for NH IFs on evidence-based recommendations about UI, and exposure to different forms of facilitation:</p> <ul style="list-style-type: none"> Type A IFs were trained in a pragmatic, quality improvement-based facilitation intervention (use of the JBI PACES) Type B IFs were trained in an enabling, critical social science-based approach, with an emphasis on inquiry, reflection and emancipatory action 	<p>IF compliance with four recommendations related to screening, assessment care planning and specialist referral</p>	<ul style="list-style-type: none"> This paper only reported outcomes related to Type A intervention Delays in introducing PACES led to a lack of baseline audit data resulting in a lack of comparative data. Qualitative data suggested IFs benefited from comparing audit data across sites
<p>Hödl et al. (2019)⁴⁰</p> <p>To measure the effectiveness of 29 evidence-based nursing recommendations regarding the conservative management of UI in Austrian NHs</p>	<p>n=381 residents randomised (100% female, mean age 85 years (IG) and 82.3 years (CG), 54.2% (IG) and 38.2% (CG) had dementia (p=0.002), 71.3% (IG) and 66.7% (CG) had UI (p=0.018))</p> <p>n=12 NHs in Austria</p>	<p>Set of 29 recommendations for conservative UI management were discussed with the NH managers and printed material was provided</p> <p>Supplementary documents (e.g., posters with abridged versions of the UI management recommendations, bladder diaries and questionnaires about QOL regarding UI) were provided</p> <p>IG: conservative UI management strategies for 12 weeks (n=216)</p> <p>CG: usual care for 12 weeks (n=165)</p>	<p>Number of daily UI events, number of UI diagnoses and use of nursing interventions were measured at baseline, 6 weeks and 12 weeks per a resident questionnaire</p>	<ul style="list-style-type: none"> IG had lower risk of daily UI (p=0.02), were less likely to receive absorbent products (p=0.01), and were five times more likely to receive recommended interventions (p=0.00) than CG

Aim(s)	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Kohler et al. (2018)⁴¹</p> <p>To explore the effects of an educational program and subsequent nursing case conferences on UI and QOL in NH residents with dementia and incontinence</p>	<p>n=140 residents randomised (79% female, mean age 85.17 years, cognitive performance scale mean score 4.06 (moderate impairment), ADL index mean 12.69) n=7 NHs in Switzerland</p>	<p>RNs, health assistants and nursing aides caring for residents received a 4-hour dementia and incontinence training session comprising presentations, group work and discussions. Staff then participated in 1-hour case conferences about persons with incontinence and dementia, occurring 2 weeks after the education session then monthly</p> <ul style="list-style-type: none"> Cluster 1: Intervention for 13 months (n=39) Cluster 2: Intervention for 11 months (n=24) Cluster 3: Intervention for 9 months (n=37) Cluster 4: Intervention for 7 months (n=40) 	<p>UI (urinary leakage via pad test) and QOL (via QualiDem) were measured at baseline, 2 and 6 months after the intervention in each cluster, and at the end of the study</p>	<ul style="list-style-type: none"> Decreases in UI by the end of the study for all clusters, significant only for Cluster 2 (T1-T3 p=0.036; T2-T3 p=0.031). QOL increased in the total sample in all categories except 'having something to do' and 'social relations', with the increase being significant for 'restless tense behaviour' (p=0.049) and 'positive self-image' (p=0.009).
<p>Lai & Wan (2017)⁴²</p> <p>To examine whether the effect of the use of PV by NH staff to manage UI among residents in NHs can be sustained for a period of 6 months</p>	<p>n=52 residents randomised (63.5% female, mean age 85.4 years, mean C-MMSE score 13.2, 32.7% severely dependent, 24.9% totally dependent) n=5 NHs in Hong Kong</p>	<p>Residents in the IG were prompted to void every 2 or 2.5 hours from 7:00am to 7:00pm for 7 days a week for 6 months</p> <p>IG: PV for 6 months (n=26)</p> <p>CG: usual care for 6 months (n=26)</p>	<p>Wet episodes per day, incontinence rate per day, self-initiated toileting per day, and total continence toileting per day, measured at baseline, 3 months post-intervention (T1), and 6 months post-intervention (T2)</p>	<ul style="list-style-type: none"> Significant differences between the two groups in wet episodes per day (p=0.001), incontinence rate per day (p=0.001), and total continence toileting per day (p≤0.001) at 6 months' post-intervention, with positive results in the IG Decrease of 9.1% in the incontinence rate in the IG
<p>Schnelle et al. (2010)⁴³</p> <p>To evaluate effects of a multicomponent intervention on FI and UI outcome</p>	<p>n=112 residents (IG 84% female, CG 81% female, mean age 85.8 years (IG) and 86.1 (CG)) n=6 NHs in the US</p>	<p>Residents were offered toileting assistance, exercise, and choice of food and fluid snacks every 2 hours for 8 hours per day over 12 weeks</p> <p>IG: multicomponent intervention for 12 weeks (n=58)</p> <p>CG: usual care for 12 weeks (n=54)</p>	<p>Frequency of UI and FI and rate of appropriate toileting as determined by direct checks from research staff. Anorectal assessments were completed on a subset of 29 residents</p>	<ul style="list-style-type: none"> Significant improvement in frequency of UI (p=0.049) Significant improvement in frequency of appropriate toileting (p=0.000) UI participants with higher MMSE scores (less cognitive impairment) responded less well to treatment (p=0.05)

Aim(s)	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Schnelle et al. (2013)⁴⁴ To evaluate an intervention to improve staff offers of choice to NH residents during morning care</p>	<p>n=138 residents randomised (76.3% female, mean age 80.5 years, 74% white, mean MMSE score 15.4, mean MDS-ADL Dependency Score 17.0) n=4 NHs in the US</p>	<p>Nursing aides caring for residents received 12 weekly video-based training sessions on how to offer choice in morning care (when get out of bed, when to get dressed/what to wear, incontinence care [changing and/or toileting], where to dine). Training sessions were followed by weekly feedback about how often choice was being provided based on standardised observations of care conducted weekly by research staff n=37 residents received the intervention immediately and completed the study n=101 residents acted as a control then received the intervention after a 12-week delay and completed the study</p>	<p>Total number of instances of "choice offered" for each participant within each morning care area and total time to provide care in the targeted areas during each observation period. Measured through standardised observations during a minimum of 4 consecutive morning hours/participant/week for 12 weeks of baseline and 12 weeks of intervention</p>	<ul style="list-style-type: none"> Significant increases in frequency that choice was offered for three care areas from baseline to intervention: out of bed, 21% to 33% (p<0.001); dressing, 20% to 32% (p<0.001); incontinence care, 18% to 23%, (p<0.014). Dining was not significant Significant increase in amount of staff time to provide care from baseline to intervention (8.01±9.0 to 9.68±9.9 minutes/person, p<0.001) (i.e., time taken to provide care was increased by the intervention)
<p>Seers et al. (2018)⁴⁵ To extend knowledge of facilitation as a process for getting research evidence into practice by testing the effectiveness of and evaluating the contribution two different models of facilitation can make to implementing evidence-based urinary continence recommendations into practice</p>	<p>n=24 long-term nursing care sites in England, Sweden, Netherlands, Republic of Ireland randomised to one of three arms Staff: one IF chosen per facility for study arms 2 and 3 (n=16) Residents: all facilities housed residents aged ≥60 years with documented UI</p>	<p>Arm 1: Lecture about four continence recommendations: <ul style="list-style-type: none"> Screen for incontinence Conduct a detailed continence assessment Design an individualised treatment plan Refer to a specialist if needed (n=8 facilities) Arm 2: per Arm 1 plus: <ul style="list-style-type: none"> 12-month goal-focused facilitation programme (n=8 facilities) Arm 3: per Arm 1 plus: <ul style="list-style-type: none"> 24-month facilitation programme with stakeholder empowerment and strategies to overcome obstacles to use of research in practice (n=8 facilities) </p>	<p>Documented percentage compliance with continence recommendations measured at baseline, 6, 12, 18 and 24 months after intervention</p>	<ul style="list-style-type: none"> No significant differences in the documented compliance with continence recommendations between study arms

Aim(s)	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Suzuki et al. (2019)⁴⁶ To determine whether USAPV care is more efficacious than conventional PV (CPV) care for managing UI in NHs</p>	<p>n=80 residents randomised (77% female, mean age 85 years, mean care-needs levels 3.5 (USAPV) and 3.2 (CPV) (moderate-high dependency)) n=13 NHs in Japan</p>	<p>n=45 residents received USAPV intervention (bladder urine volume monitored by an ultrasound device and staff prompted them to void accordingly) n=35 residents received standard PV (residents asked by staff every 2-3 hours whether they had a desire to void and were prompted to void when response was yes)</p>	<p>Primary outcomes: • Change in daytime urine loss and QOL assessed using the EQ-5D from baseline to end of intervention Secondary outcomes: • Scores of mental state, depression, physical function, and level of motivation assessed using the EQ-5D, MMSE, GDS, Barthel index, and vitality index, respectively • Change in caregivers' QOL assessed with the SF-12v223 and mental stress for care burden with the visual analogue scale (VAS)</p>	<ul style="list-style-type: none"> • Change in daytime urine loss statistically greater for USAPV (median -80.0g) than CPV (median -9.0g; p=0.018) group • Proportion of residents whose daytime urine loss decreased by >25% was 51% and 26% in the USAPV and CPV group, respectively (p=0.020) • No significant changes in QOL for either group • Care burden was unchanged in the USAPV group (p=0.59) but significantly worsened in the CPV group (p=0.010) after the intervention
<p>Tak et al. (2012)⁴⁷ To compare a group-based behavioural exercise program to prevent or reduce UI, with usual care. The exercise program aimed to improve functional performance of pelvic floor muscle, bladder and physical performance of women living in homes for the elderly</p>	<p>n=192 female residents randomised (100% female, mean age 84.6 years (IG) and 84.7 (CG), with or without UI, with Cognitive Screening Test score >9.6 (good cognition) and able to use the toilet independently) n=20 NHs in the Netherlands</p>	<p>Group-based behavioural exercise program comprising weekly group exercises and independent homework exercises for 6 months IG: exercise program for 6 months (n=85) CG: usual care for 6 months (n=70)</p>	<p>Presence or absence of UI, frequency of episodes (3-day bladder diary), Physical Performance Test, SF-12 physical, SF-12 mental and I-QOL measured at baseline, 3 and 6 months</p>	<ul style="list-style-type: none"> • Physical performance (PPT) significantly improved in the IG compared to a decline in the CG (p≤0.001) • No significant differences between groups on any other outcome measures
<p>Vinsnes et al. (2012)⁴⁸ To investigate if an individualised training program designed to improve ADL and physical capacity among residents in NHs has any impact on UI</p>	<p>n=98 residents randomised (75.5% female, mean age 84.3 years, mean MMSE score 12.5, and in need of assistance with at least one ADL) n=24 NHs in Sweden, Norway and Denmark</p>	<p>Personal/individualised treatment goals were elicited for each participant. Each participant was then involved in a training program including accommodated physical activity and ADL training for 3 months IG: training program for 3 months (n=35) CG: usual care for 3 months (n=33)</p>	<p>UI as measured by a 24-hour pad-weighing test, measured at baseline and 3 months post-intervention</p>	<ul style="list-style-type: none"> • Amount of leakage significantly decreased in the IG after adjusting for baseline leakage, age, sex and functional status (p=0.03)

Aim(s)	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Wijk et al. (2018)⁴⁹</p> <p>To operationalise, assess and evaluate the feasibility and preliminary effects of implementing a person-centred approach to incontinence care for older adults with cognitive decline in NHs in Sweden</p>	<p>n=79 residents randomised (59.9% female, mean age 83.9 years, MMSE score <23)</p> <p>n=2 NHs in Sweden</p>	<p>Staff (n=24) caring for residents received a 10-month training program about how to encounter and interview residents from a person-centred approach and create person-centred care plans, and received training in incontinence assessment, planning, caring actions and outcome follow-up. Staff used this training to elicit resident narratives and incorporate these narratives into resident care plans</p> <p>IG: residents receiving care from trained staff facilitators (n=2 NHs)</p> <p>CG: residents receiving usual care (n=1 NH)</p>	<p>Process outcome:</p> <ul style="list-style-type: none"> Number of assessments conducted for incontinence management <p>Impact outcomes:</p> <ul style="list-style-type: none"> QOL (QUALID scale), basal assessment of incontinence, incontinence actions taken, personally chosen incontinence aids <p>Measured at baseline, during implementation, immediately after end of implementation, and 6 months after end of implementation</p>	<ul style="list-style-type: none"> Non-significant increase in residents' QOL in IG compared to baseline and CG Positive effect on the number of UI assessments conducted in the intervention NHs at baseline and follow up 1 (p≤0.05) Number of person-centred caring actions (e.g., toilet assistance) was significantly higher during and 6 months after implementation of the person-centred approach (p=0.019)

to identify whether different forms of facilitation were influential within very diverse contextual conditions.

In the US, Schnelle et al.⁴⁴ conducted a controlled trial to evaluate the effects of educating nursing home staff about offering residents greater choice about toileting assistance, exercise and choice of food and fluid snacks. They were encouraged to offer this choice every 2 hours for 8 hours per day over 3 months. At 3 months, the researchers reported significant increases in the number of time residents received choice-related toileting (p=0.014). However, they also noted a significant increase in the amount of staff time to provide care (p≤0.001).

Gencbas et al.³⁸ evaluated the effects of educating nursing home staff on the use of a standardised nursing language to identify residents with UI and to design targeted nursing interventions in five nursing homes in Turkey. The cluster RCT conducted over 12 weeks was evaluated with respect to residents' continence status and QOL. The researchers reported a between-group statistically significant improvement in the use of the standardised nursing language, residents' QOL and symptom severity.

All four quasi-experimental studies that examined the effects of educating nursing home staff about UI were conducted in the US⁵¹⁻⁵⁴. They included a 6-week pre/post-study by Mathis et al.⁵² involving 166 nursing home staff from six US nursing homes. The education focussed on the types and treatment of incontinence and was delivered face-to-face. The researchers reported significant improvements in post-education staff knowledge about types of UI (p≤0.001 for all domains) and improved attitudes about UI (p≤0.001), but no significant change in staff knowledge about assessment or treatment.

By contrast, Ehlman et al.⁵¹ reported significant improvements in nursing home staff members' attitudes about UI following nine in-service programs designed to teach 107 staff from four skilled nursing facilities in the US how to use a bladder scanner. This was augmented with a DVD on the same topic, a refresher in-service 12 weeks later, and supplementary information and incentives. Post-intervention improvements were also noted for knowledge about using a bladder scanner, but not for the belief that UI is a normal part of ageing.

Rahman et al.^{53,54} compared a 5-month course about PV delivered by teleconference with an 8-month course. The researchers reported a significant increase in nursing home staff knowledge compared to baseline (p≤0.05 and p≤0.001) and that varying the number and duration of teleconference sessions did not produce any significant effects. The researchers also evaluated the effects of six 40-minute webinars on the same topic⁵⁴. Based on nursing home documentation, 100% of participating homes implemented toileting assistance following the education and 83% reported that they planned to continue the intervention.

Multi-component interventions

Multi-component interventions that combine

Table 4. Characteristics of included studies: quasi-experimental studies

Aim	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Arnold et al. (2016)⁵⁰</p> <p>To assess if conservative therapy can reduce urinary leakage and pad usage and improve QOL in elderly incontinent women living in a rest home setting and, if so, at what additional cost</p>	<p>n=68 residents (100% female, mean age 85.3, mean MMSE score 25.6)</p> <p>n=26 NHs in New Zealand</p>	<p>Residents received tailored, conservative continence treatment from a specialist continence nurse advisor for 12 weeks.</p> <p>Interventions included pelvic floor muscle exercises and bladder retraining with a focus on deferment strategies, fluid intake, optimising bowel function and increased physical activity, and lifestyle changes. PV, techniques to improve bladder emptying, prescription of anticholinergics and oestrogen cream were also recommended to some residents</p>	<p>Primary outcomes: Bladder function (via bladder diary, pad weigh tests, pad usage), and QOL and activities via questionnaires (FIM); Functional Impairment Measure; EQ-5D; EuroQoL 5-dimension score; ICIQ-SF; Incontinence-Short Form) recorded at baseline and after 12 weeks</p> <p>Secondary outcomes: Pad cost, carers' cost for continence care, additional laundry costs and total costs</p>	<ul style="list-style-type: none"> • Mean urine loss fell significantly by 60ml per day (p=0.024) • Significant reduction in pad changes per week by a mean of 4.3 (p<0.0005) • ICIQ-SF improved statistically by 2.84 (p<0.0005) but no significant improvement in any other QOL or activities scales • Pads used per week reduced by a mean of 4.3 with a significant reduction of pad costs per week (mean NZ\$2.43/16%) • No significant different in carers' cost or additional laundry costs • Total cost of providing the intervention for 12 weeks (NZ\$247.75 minus cost-savings NZ\$12.48), was NZ\$235.27 per participant
<p>Ehlman et al. (2012)⁵¹</p> <p>To explore differences in attitude and knowledge related to UI among nursing personnel who provide care for NH residents, and examine staff attitude and knowledge about UI</p>	<p>n=107 NH staff (15% registered nurses, 54.2% licensed practical nurse (LPN), 30.8% CNA)</p> <p>n=4 skilled nursing facilities in the US</p>	<p>48 aged care staff completed in-service education about UI and about the use of a bladder ultrasound machine. Facilities had a bladder ultrasound machine made available, and staff completed a pre- and post-intervention surveys</p>	<p>Staff attitudes and knowledge regarding incontinence</p>	<ul style="list-style-type: none"> • Before the intervention, there was a significant association between position and belief that bladder disorders are a normal part of ageing (more CNAs than LPNs believed this [p<0.003] as did more CNAs than RNs [p=0.003]) • Study intervention resulted in a significant positive change in attitude toward UI (p<0.001)
<p>Mathis et al. (2013)⁵²</p> <p>To determine: staff knowledge about the types of UI before and after participating in the Bladder Buzz program; staff attitudes about UI before and after participating in the program; and staff knowledge about the treatment and management of UI before and after participating in the program</p>	<p>n=166 NH staff (14% registered nurses, 23% LPN, 39% CNA and 23% non-nursing staff)</p> <p>n=6 NHs in the US</p>	<p>38 NH staff members completed a 6-week educational intervention on the types and treatment of UI and attitudes toward UI, and completed pre- and post-intervention surveys</p>	<p>Staff knowledge about UI, attitudes toward UI, and treatment and assessment of UI assessed via a researcher-developed survey before and after the intervention</p>	<ul style="list-style-type: none"> • Changes in knowledge about the types of UI were significant for stress (p<0.001), functional (p<0.003), and overflow (p<0.00) UI • Changes in attitudes toward UI were significant (p<0.000) • Changes in staff knowledge of the assessment and treatment of UI were not significant

Aim	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Rahman et al. (2012)⁵³</p> <p>To determine whether a distance coaching course on improving NH incontinence care could be replicated and brought to scale with a larger group of NHs without sacrificing outcomes</p>	<p>Supervisors and staff (not further described or quantified) from NHs in the US</p>	<p>Course 1: PV education delivered to 14 staff via 8 monthly teleconferences of 60-90 minutes each</p> <p>Course 2: PV education delivered to 34 staff via 5-monthly teleconferences and one 2-month follow-up of 40 minutes each</p>	<p>Number of staff participating in course, implementation of PV intervention, effectiveness of training in increasing knowledge, training program preferences, and course costs</p>	<ul style="list-style-type: none"> No significant differences in participation, intervention implementation, knowledge increase or training preferences were found between groups Both groups showed a significant increase in knowledge compared to baseline (Course 1 $p \leq 0.05$, Course 2 $p \leq 0.0001$). The cost of Course 1 was US\$1894/facility and the cost of Course 2 was S\$514/participating facility
<p>Rahman et al. (2014)⁵⁴</p> <p>To administer and evaluate a webinar course that provided extended instruction to help NHs implement toileting trials in accordance with recommended procedures</p>	<p>Staff members (nurses, DONs, MDS coordinators, staff educators, nursing aides) (not further quantified or described) from n=7 NHs in the US</p>	<p>Six 40-minute webinars comprising an online course to assist NH staff to implement toileting trials were delivered to 16 staff over 9 weeks. Between webinars, staff were asked to complete implementation "assignments":</p> <ul style="list-style-type: none"> Interview a sample of incontinent residents to assess their preferences for incontinence care Conduct a 3-day toileting trial with these residents Re-interview the residents following the trial Summarise the results from this procedure <p>Facility "champions" attended all webinars and oversaw implementation of assignments (n=7)</p>	<p>Whether the course prompted NH staff to adopt the best-practice toileting trial protocol (evidenced by documentation of new toileting trial assessments being completed for incontinent residents)</p> <p>Evaluation of the course itself (whether staff participants preferred this course to more traditional continuing education models) via course and pilot-test evaluation surveys completed by staff and champions</p>	<p>Primary evaluation results:</p> <ul style="list-style-type: none"> 100% of champions reported that their staff had completed all three toileting trial steps (i.e., interview, trial and analysis) with at least some residents 50% of champions reported that, in their opinion, assessed residents were drier and that more incontinent residents had individualised toileting plans as a result of course participation 83% of champions said that their facility planned to continue or expand use of all three standardised continence assessment forms that were used in the toileting trials <p>Secondary evaluation results:</p> <ul style="list-style-type: none"> 83% of champions and 75% of staff said they preferred the webinar course to other course designs
<p>Suzuki et al. (2016)⁵⁵</p> <p>To assess the efficacy and feasibility of USAPV for the management of UI in NHs</p>	<p>n=77 residents (76.6% female, mean age 84.1, primarily dependent [53.2% classified as care needs level 4 or 5]) n=6 NHs in Japan</p>	<p>Residents received a PV intervention for 1 week, followed by USAPV intervention for 12 weeks (residents were prompted to void on the toilet when monitored urine volume reached close to that optimal volume that had calculated for each resident)</p>	<p>Primary outcome:</p> <ul style="list-style-type: none"> Change in absorbent cost from baseline to the end of the study <p>Secondary outcomes:</p> <ul style="list-style-type: none"> ADLs of participants assessed using the vitality index Feasibility of the intervention assessed via care worker QOL using SF-36 	<ul style="list-style-type: none"> Absorbent cost was reduced in 40 of the 77 participants (51.9%) Being pad-free was attained by two participants (2.9%) at 4 weeks, two at 8 weeks and three (3.9%) at 12 weeks, respectively Change in total vitality index score from 6.96 to 6.76 was not significant ($p=0.236$) Two subscales of the SF-36 Health Survey, role emotional ($p=0.020$) and mental health ($p=0.007$), improved significantly after intervention

Aim	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Yu et al. (2014)⁵⁶</p> <p>To explore the effects of introducing a telemonitoring and care planning system for urinary continence assessment in a NH and adherence by care staff to urinary continence care plans</p>	<p>n=31 residents (78% female, mean age 81±8) n=1 NH in Australia</p>	<p>Staff who care for residents were trained in the use of video telemonitoring for continence assessment over a 12-week trial</p>	<ul style="list-style-type: none"> Weight of urine voided into the continence aid (episode of UI void) Number of prescribed toileting events in the care plans Number of actual toileting events Number of successful toileting events Secondary measure of the rate of successful toileting Adherence to care plans by staff <p>Measured at baseline and post-intervention</p>	<ul style="list-style-type: none"> Significantly reduced volume of urine voided into continence aids (p=0.015) Significantly reduced number of prescribed toileting visits (p=0.015) Significantly increased number of actual toilet visits (p<0.001) Significantly increased number of successful toileting events (p=0.011) Significantly increased adherence to urinary continence care plans by staff (p=0.033)

Table 5. Characteristics of included studies: retrospective cohort studies

Aim(s)	Setting/baseline sample	Intervention(s)	Outcomes measured	Findings
<p>Moga et al. (2013)⁵⁷</p> <p>To evaluate risks and benefits of bladder antimuscarinics (BAMs) among elderly NH residents</p>	<p>Retrospective cohort study of older adults (65 years and older) admitted for long-term care between 1 October 2002 and 30 September 2009 in Veterans Affairs (VA) community living centres (NHs)</p>	<p>Retrospective cohort study. Through a new user design, BAM users were compared with propensity score (PS)-matched non-users and the impact of BAM initiation was evaluated</p> <p>A new user was an individual whose first dispensing of a BAM (oxybutynin chloride, tolterodine, darifenacin, solifenacin, trospium, hyoscyamine, dicyclomine, orflavoxate; immediate-release [IR] or extended-release [ER] formulations) occurred more than 3 days after being admitted to a VA NH and who did not have any BAM prescription within 1 year before this first prescription date</p>	<p>Treatment benefits were measured through change in urinary continence status and QOL, whereas fracture risk and decline in cognitive performance were the potential harmful effects studied</p>	<p>BAM initiation was significantly associated with improvement in UI (e.g., from frequently incontinent to occasionally incontinent or from occasionally incontinent to usually continent) (OR=1.34, 95% CI 1.13-1.60). The addition of time-varying covariates did not significantly change the estimate (OR=1.27, 95% CI 1.07-1.5).</p> <p>At 90 days after BAM initiation, 60 new users (16.35%) showed improvement (as compared with 13.02% for non-users). After 180 days, 93 new users (25.34%) had a positive outcome in terms of UI improvement compared with 20.34% in the non-users group.</p> <p>The NNT was 32 (95% CI 17-125), indicating that 32 patients must be treated for 90 days to obtain improvement in UI in one patient. The risk of fractures was significantly increased in new users compared to non-users (hip fracture: hazard ratio [HR]=3.67, 95% CI 1.46-9.34; 'any' fracture: HR=2.64, 95% CI 1.37-5.10)</p>

physical activity with other interventions can reduce UI in nursing homes according to the findings of a systematic review²⁹. In addition, two RCTs demonstrated the effectiveness of multi-component interventions that combine physical activity with other interventions. A trial by Schnelle et al.⁴³ evaluated exercise in combination with toileting assistance and choice of food and fluid snacks every 2 hours for 8 hours per day over 12 weeks. Compared to controls (n=60), residents (n=65) who received this multi-component intervention had a significant reduction in the frequency of UI ($p=0.049$) and an improvement in the frequency of appropriate toileting ($p\leq 0.001$). Azizi et al.³⁵ compared the effectiveness of an 8-week multi-component program (light strengthening exercises, pelvic floor exercises, and an education program about UI with usual care). Residents (n=30) who participated in the program showed a significant decrease in UI severity ($p\leq 0.001$).

An additional three publications described the effect of multi-component interventions, albeit specifically focussing on the role of specialist healthcare professionals. They were a systematic review²⁸, a cluster RCT³⁶ and a quasi-experimental study⁵⁰. The systematic review reported on four studies in which a clinical nurse specialist (CNS) facilitated evidence-based protocols, collaborated with staff, and provided staff education, consultation and direct care to residents for 6 months post-admission²⁸. Residents in nursing homes who received these interventions showed reduced UI or slowed decline in UI²⁸. This finding is further supported by a subsequent quasi-experimental study involving 68 nursing home residents who received tailored, conservative continence treatment from a continence nurse advisor for 12 weeks⁵⁰. A variety of strategies were implemented for each resident. After implementation, mean urine loss fell significantly by 60ml per day ($p=0.024$) and a significant reduction in pad changes per week by a mean of 4.3 ($p\leq 0.0005$) was observed. Incontinence-related QOL scores improved significantly ($p\leq 0.005$) but there was no significant improvement in any other QOL or activities scale measured.

In the cluster RCT, Boorsma et al.³⁶ compared a multidisciplinary integrated care program with usual care in 10 nursing homes in the Netherlands. The program comprised assessments, individualised care plans, discussions between the family physician, resident and their family, and multidisciplinary staff meetings to discuss residents with complex needs. Each component of the program was repeated quarterly for the duration of the study. Compared with the facilities that provided usual care, intervention facilities had a significantly higher sum score of 32 quality of care indicators and significantly higher scores for 11 of 32 indicators of good care, including UI.

DISCUSSION

This scoping review was undertaken to map and summarise evidence about interventions for the management of UI in nursing homes. We identified

several different intervention types and grouped them according to whether they represented a toileting assistance program, an exercise program, drug therapy, a technology-based intervention, an education program, or a multi-component intervention.

We identified six publications about toileting assistance programs, namely PV — three systematic reviews^{29,31,32}, two RCTs^{42,46} and one quasi-experimental study⁵⁵. The latter three studies provide recent evidence to support the use of technology to augment assessment processes and for the implementation and sustainability of PV by usual care staff. The use of ultrasound technology to identify when nursing home residents need to void is a relatively new innovation for the management of incontinence in nursing homes, as is the use of sensor alarms. The technology is based on the observation that some nursing home residents' have predictable voiding patterns and that, by identifying the pattern, staff will be better placed to provide targeted and timely assistance. As the technology can provide more accurate and objective information about a person's actual bladder function, they can augment current assessment processes and thereby minimise trial and error. However, important questions remain about the transferability of the findings to different cultural contexts, staff training needs and costs as well as residents' acceptance of the technology. It is important to understand the relationship between the effectiveness of technological interventions in nursing homes, the provision of staff or carer training and support, and the process of fidelity to the intervention.

Recent evidence also points to the need to optimise nursing home residents' functional skills and mobility¹¹. Despite this, only two studies specifically described an evaluation of an exercise program, and the findings were inconsistent. The best evidence is for multi-component interventions that combine physical activity programs with systematic efforts to improve residents' choice, food and fluid intake, and access to toileting assistance. Again, studies point to the value of engaging advanced practice registered nurses to facilitate this approach.

This scoping review found limited evidence about the use of drug therapies for the management of UI in nursing homes. Based on two systematic reviews and a cohort study, the benefits of using anticholinergics are outweighed by the risks of adverse effects. It is also clear that, although acceptable, transcutaneous posterior tibial nerve stimulation is unlikely to confer a reduction in UI in nursing homes with a high proportion of older frail residents with reduced cognitive capacity.

Staff education was the focus of 11 publications. Prior research points to gaps in undergraduate education programs for healthcare practitioners in the UK⁵⁸ and in the US⁵⁹. A recent review of the effects of UI education for nurses and nursing assistants' knowledge, attitudes and self-reported practices found education improved knowledge, but the most effective forms of education that affects practice and patient outcomes are not known³⁰.

Nursing home staff should be educated on best practice guidelines. The findings from a large cluster RCT reported by Harvey et al.³⁹ and Seers et al.⁴⁵ highlight the complexities of implementing and evaluating guidelines in nursing homes, even when it is facilitated. The researchers found no significant differences between groups in the documented percentage compliance with the continence recommendations. There could be several reasons for this finding, including limitations of the guidelines themselves. Many guidelines about incontinence are underpinned by the medical goal of diagnosis and treatment. We argue that guidelines and education programs about continence care should place equal value on helping people adjust to changes in bodily function that affect their identity, autonomy, control and independence.

Contemporary forms of education about care in health and social care systems promote person-centred care facilitated through a shared-decision making process⁶⁰⁻⁶². This highlights the importance of the intervention implemented and evaluated by Wijk et al.⁴⁹ where nursing home staff were educated on how to elicit information from residents about their personal identity and preferences and to then use this information to design individualised continence care plans. It should be noted that introducing a person-centred approach to continence care takes time and commitment and may require a culture change for some nursing homes.

Given the high proportion of residents with a diagnosis of dementia in nursing homes, education programs for staff should also address the prevention and management of incontinence in people with dementia. The education program described by Kohler et al.⁴¹ is an important exemplar. The program addressed dementia as well as incontinence and was augmented with case conferences for staff to discuss problematic and challenging situations concerning specific nursing home residents. Our analysis of education interventions also points to the need for direct hands-on education combined with on-site support and competency-based learning approaches, particularly when led by specialist healthcare professionals.

Two studies provide evidence that organisational structures, staff resources, staffing models and care processes impact on nursing home residents' risk of incontinence^{63,64}. Work environment attributes such as team cohesiveness, consistent assignment and staff cohesion affect residents' urinary continence status⁶³. Within this context, higher registered nurse to patient ratios is an important consideration. A longitudinal correlational study on the impact of organisational factors on the continence status of patients in Korean long-term care hospitals found higher registered nurse to patient ratios was significantly associated with better resident continence status⁶⁴. These key findings highlight the important role that registered nurses and nursing home leaders have in creating a therapeutic physical and socio-cultural environment that facilitates continence.

Based on one study and a systematic review, we also recommend multicomponent interventions delivered by registered nurses with advanced practice skills^{28,50} and for multidisciplinary integrated care³⁶. Ideally, the multicomponent intervention should consist of assessments, individualised care plans, discussions between the family physician, residents and their families, and multidisciplinary staff meetings to discuss residents with complex needs.

Strengths and limitations

A weakness of this scoping review was that we did not develop and register an a priori protocol and thus the review does not fully adhere to the *JBI reviewers' manual 2015 methodology for JBI scoping reviews*²⁵, nor the *PRISMA-ScR guidelines*²⁶. At the same time, a key strength of this scoping review is that the broad nature of the design yielded evidence about several types of interventions for the management of UI in nursing homes.

The review was limited to research with quantitative findings. As such, it focussed on questions of effectiveness rather than context. Further reviews are required to summarise qualitative research about the psychological and sociological aspects of UI in nursing homes.

Finally, nursing homes vary considerably, both within and between countries. Therefore, interventions that are effective in one context may not be effective in another. Aligned with this consideration is the cultural bias in the body of evidence. Most of the research identified was conducted in Western countries and none from lower to middle income countries. Of the 30 studies identified, 18 were conducted in nursing homes in Western countries. Hence, the transferability to non-Western and lower to middle income countries should be considered. Further research is also needed to understand whether existing UI management strategies are able to meet the specific needs of residents from diverse backgrounds, such residents from culturally and linguistically diverse (CALD) communities, First Nations residents, and residents who identify as lesbian, gay, bisexual, transgender and/or intersex (LGBTIQ+).

Finally, while each study was appraised, studies were not excluded from the review based on their quality. Moreover, findings were reported descriptively and were not subject to rigorous assessment, taking potential biases into account. In several of the studies unblinding may have introduced significant bias; however, given the nature of some of the interventions (ie, personalised therapies, physical activity programs and workshops), unblinding may be acceptable in this context.

CONCLUSION

UI continues to be a pervasive problem that disproportionately affects people in nursing homes. Interventions to maintain or improve nursing home residents' functional status and reduce their care

dependence should be prioritised. Improvements in staff knowledge, attitudes and continence care practices in nursing homes are possible, particularly when led by specialist healthcare professionals.

This scoping review suggests a multi-component approach to care offers most benefit, ie, where efforts are made to increase residents' activity level, nutritional status, hydration and opportunity to use the toilet. This approach should also be combined with direct forms of education for staff to optimise compliance. Future directions for policy should promote person-centred continence care, comprehensive multi-disciplinary assessments, and a well-educated and supported workforce. Registered nurses play a key role in creating a therapeutic physical and socio-cultural environment that facilitates continence in nursing homes.

AUTHOR CONTRIBUTIONS

All of the authors made substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data. They were all involved in drafting the manuscript or revising it critically for important intellectual content and gave final approval of the version to be published.

ACKNOWLEDGEMENTS

The researchers wish to acknowledge input to the research study by Ms Erica Wise and Dr Stephanie Garratt.

REFERENCES

- Wagg A, Bower W, Gibson W, Kirschner-Hermanns R, Hunter K, Kuchel GA, Morris V, Ostaszkievicz J, Suskind A, Suzuki M, Wyman J. Incontinence in frail older adults. In: Cardozo L, Rovner E, Wagg A, Wein A, Abrams P, editors. *Incontinence: 7th international consultation on incontinence*, vol 13. London: ICUD ICS; 2023.
- Boyington JE, Howard DL, Carter-Edwards L, Gooden KM, Erdem N, Jallah Y, Busby-Whitehead J. Differences in resident characteristics and prevalence of urinary incontinence in nursing homes in the southeastern United States. *Nurs Res* 2007;56(2):97-107.
- Jachan DE, Müller-Werdan U, Lahmann NA. Impaired mobility and urinary incontinence in nursing home residents: a multicenter study. *J Wound Ostomy Continence Nurs* 2019;46(6):524-529.
- Harrison T, Blozis S, Manning A, Dionne-Vahalik M, Mead S. Quality of care to nursing home residents with incontinence. *Geriatr Nurs* 2019;40(2):166-173.
- Mandl M, Halfens RJ, Lohrmann C. Interactions of factors and profiles of incontinent nursing home residents and hospital patients: a classification tree analysis. *J Wound Ostomy Continence Nurs* 2016;43(4):407-413.
- Mid Staffordshire NHS Foundation Trust. Report of the Mid Staffordshire NHS Foundation Trust public inquiry: executive summary. Staffordshire: Mid Staffordshire NHS Foundation Trust; 2013. Available from: <http://www.midstaffpublicinquiry.com/sites/default/files/report/Executive%20summary.pdf>
- The Royal Commission into Quality and Safety in Aged Care. Final report: care dignity and respect. Canberra: Royal Commission into Quality and Safety in Aged Care; 2021. Available from: <https://agedcare.royalcommission.gov.au/publications/final-report>
- Ostaszkievicz J, Tomlinson E, Hutchinson A. 'Dignity': a central construct in nursing home staff understandings of quality continence care. *J Clin Nurs* 2018;27:2425-2437.
- Ostaszkievicz J, Tomlinson C, Hutchinson A. Learning to accept incontinence and continence care in residential aged care facilities: family members' experiences. *Aust NZ Cont J* 2016; 22(3):14-21.
- MacDonald CD, Butler L. Silent no more: elderly women's stories of living with urinary incontinence in long-term care. *J Gerontolog Nurs* 2007;33(1):14-20.
- Li HC, Chen KM, Hsu HF. Modelling factors of urinary incontinence in institutional older adults with dementia. *J Clin Nurs* 2019;28(23-24):4504-4512.
- Watson NM, Brink CA, Zimmer JG, Mayer RD. Use of the Agency for Health Care Policy and Research Urinary Incontinence Guideline in nursing homes. *J Am Geriatr Soc* 2003;51(12):1779-1786.
- Xu D, Kane RL. Effect of urinary incontinence on older nursing home residents' self-reported quality of life. *J Am Geriatr Soc* 2013;61(9):1473-1481.
- Jerez-Roig J, Moreira FSM, da Câmara SMA, Ferreira LMBM, Lima KC. Predicting continence decline in institutionalized older people: a longitudinal analysis. *Neurourol Urodyn* 2019;38(3):958-967.
- Van Damme N, Van den Bussche K, De Meyer D, Van Hecke A, Verhaeghe S, Beeckman D. Independent risk factors for the development of skin erosion due to incontinence (incontinence-associated dermatitis category 2) in nursing home residents: results from a multivariate binary regression analysis. *Int Wound J* 2017;14(5):801-810.
- Hasegawa J, Kuzuya M, Iguchi A. Urinary incontinence and behavioural symptoms are independent risk factors for recurrent and injurious falls, respectively, among residents in long-term care facilities. *Arch Gerontol Geriatr* 2010;50(1):77-81.
- Moon S, Hong GS. Predictive factors of mortality in older adult residents of long-term care facilities. *J Nurs Res* 2020;28(2):e82.
- Frantz RA, Xakellis Jr GC, Harvey PC, Lewis AR. Implementing an incontinence management protocol in long-term care: clinical outcomes and costs. *J Gerontolog Nurs* 2003;29(8):46-53.
- Schnelle JF. Treatment of urinary incontinence in nursing home patients by prompted voiding. *J Am Geriatr Soc* 1990;38:356-360.
- Schnelle JF, Newman DR, Fogarty TE, Wallston K, Ory M. Assessment and quality control of incontinence care in long term nursing facilities. *J Am Geriatr Soc* 1991;36:165-71.
- Schnelle JF, Newman D, White M, Abbey J, Wallston KA, Fogarty T, Ory MG. Maintaining continence in nursing home residents through the application of industrial quality control. *Gerontologist* 1993;33:114-121.
- Schnelle JF, MacRae PG, Ouslander JG, Simmons SF, Nitta M. Functional incidental training, mobility performance, and incontinence care with nursing home residents. *J Am Geriatr Soc* 1995;43(12):1356-62.
- Schnelle JF, Alessi CA, Simmons SF, Al-Samarrai NR, Beck JC, Ouslander JG. Translating clinical research into practice: a randomized controlled trial of exercise and

- incontinence care with nursing home residents. *J Am Geriatr Soc* 2002;50:1476-1483.
24. Eustice S, Roe B, Paterson J. Prompted voiding for the management of urinary incontinence in adults. *Cochrane Database System Rev* 2000;(2), CDO02113.
 25. Peters M, Godfrey C, McInerney P, Soares CB, Khalil H, Parker D. The Joanna Briggs Institute reviewers' manual 2015: methodology for JBI scoping reviews. The Joanna Briggs Institute; 2015.
 26. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D, Peters MDJ, Horsley T, Weeks L, Hempel S, Akl EA, Chang C, McGowan J, Stewart L, Hartling L, Aldcroft A, Wilson MG, Garritty C, Lewin S, Straus SE. PRISMA Extension for Scoping Reviews (PRISMA-ScR): checklist and explanation. *Ann Int Med* 2018;169(7):467-473.
 27. Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, Hartmann-Boyce J, Ryan R, Shepperd S, Thomas J, Welch V, Thomson H. Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *Br Med J* 2020;368:l6890.
 28. Donald F, Martin-Misener R, Carter N, Donald EE, Kaasalainen S, Wickson-Griffiths A, Lloyd M, Akhtar-Danesh N, DiCenso A. A systematic review of the effectiveness of advanced practice nurses in long-term care. *J Adv Nurs* 2013;69(10):2148-2161.
 29. Flanagan L, Roe B, Jack B, Barrett J, Chung A, Shaw C, Williams KS. Systematic review of care intervention studies for the management of incontinence and promotion of continence in older people in care homes with urinary incontinence as the primary focus (1966-2010). *Geriatr Gerontol Int* 2012 Oct;12(4):600-11.
 30. Ostaszkiwicz J, Tomlinson E, Hunter K. The effects of education about urinary incontinence on nurses' and nursing assistants' knowledge, attitudes, continence care practices and patient outcomes. *J Wound Ostomy Continence Nurs* 2020;47(4):365-380.
 31. Roe B, Flanagan L, Jack B, Barrett J, Chung A, Shaw C, Williams K. Systematic review of the management of incontinence and promotion of continence in older people in care homes: descriptive studies with urinary incontinence as primary focus. *J Adv Nurs* 2011;67(2):228-250.
 32. Roe B, Flanagan L, Maden M. Systematic review of systematic reviews for the management of urinary incontinence and promotion of continence using conservative behavioural approaches in older people in care homes. *J Adv Nurs* 2015;71(7):1464-1483.
 33. Samuelsson E, Odeberg J, Stenzelius K, Molander U, Hammarström M, Franzen K, Andersson G, Midlöv P. Effect of pharmacological treatment for urinary incontinence in the elderly and frail elderly: a systematic review. *Geriatr Gerontol Int* 2015;15(5):521-534.
 34. Stenzelius K, Molander U, Odeberg J, Hammarström M, Franzen K, Midlöv P, Samuelsson E, Andersson G. The effect of conservative treatment of urinary incontinence among older and frail older people: a systematic review. *Age Ageing* 2015;44(5):736-744.
 35. Azizi M, Azadi A, Ottaghi M. The effect of a self-care programme on urinary incontinence and self-esteem in elderly men dwelling in nursing homes in Iran. *Aging Male* 2020;23(5):687-693.
 36. Boorsma M, Frijters DH, Knol DL, Ribbe ME, Nijpels G, van Hout HP. Effects of multidisciplinary integrated care on quality of care in residential care facilities for elderly people: a cluster randomized trial. *Can Med Assoc J* 2011;183(11):E724-E732.
 37. Booth J, Aucott L, Cotton S, Davis B, Fenocchi L, Goodman C, Hagen S, Harari D, Lawrence M, Lowndes A, Macaulay L, MacLennan G, Mason H, McClurg D, Norrie J, Norton C, O'Dolan C, Skelton D, Surr C, Treweek S. Tibial nerve stimulation compared with sham to reduce incontinence in care home residents: ELECTRIC RCT. *Health Technol Assess* 2021;25(41):1-110.
 38. Gencbas D, Bebis H, Cicek H. Evaluation of the efficiency of the nursing care plan applied using NANDA, NOC, and NIC linkages to elderly women with incontinence living in a nursing home: a randomized controlled study. *Int J Nurs Know* 2018;29(4):217-226.
 39. Harvey G, Kitson A, Munn Z. Promoting continence in nursing homes in four European countries: the use of PACES as a mechanism for improving the uptake of evidence-based recommendations. *Int J Evid-based Health* 2012;10(4):388-396.
 40. Hödl M, Halfens RJG, Lohrmann C. Effectiveness of conservative urinary incontinence management among female nursing home residents: a cluster RCT. *Arch Gerontol Geriatr* 2019;81:245-251.
 41. Kohler M, Schwarz J, Burgstaller M, Saxer S. Incontinence in nursing home residents with dementia: influence of an educational program and nursing case conferences. *Gerontologie und Geriatrie* 2018;51(1):48-53.
 42. Lai CKY, Wan X. Using prompted voiding to manage urinary incontinence in nursing homes: can it be sustained? *J Am Med Dir Assoc* 2017;18(6):509-514.
 43. Schnelle JF, Leung FW, Rao, SS, Beuscher L, Keeler E, Clift JW, Simmons S. A controlled trial of an intervention to improve urinary and faecal incontinence and constipation. *J Am Geriatr Soc* 2010;58(8):1504-1511.
 44. Schnelle JF, Rahman A, Durkin DW, Beuscher L, Choi L, Simmons SF. A controlled trial of an intervention to increase resident choice in long term care. *J Am Med Dir Assoc* 2013;14(5):345-351.
 45. Seers K, Rycroft-Malone J, Cox K, Crichton N, Edwards RT, Eldh AC, Estabrooks CA, Harvey G, Hawkes C, Jones C, Kitson A, McCormack B, McMullan C, Mockford C, Niessen, T, Slater P, Titchen A, van der Zijpp T, Wallin L. Facilitating Implementation of Research Evidence (FIRE): an international cluster randomised controlled trial to evaluate two models of facilitation informed by the Promoting Action on Research Implementation in Health Services (PARiHS) framework. *Implement Sci* 2018;13(1):137.
 46. Suzuki M, Miyazaki H, Kamei J, Yoshida M, Taniguchi T, Nishimura K, Igawa Y, Sanada H, Homma Y. Ultrasound-assisted prompted voiding care for managing urinary incontinence in nursing homes: a randomized clinical trial. *Neurourol Urodyn* 2019;38(2):757-763.
 47. Tak EC, van Hespden A, van Dommelen P, Hopman-Rock M. Does improved functional performance help to reduce urinary incontinence in institutionalized older women? a multicenter randomized clinical trial. *BMC Geriatr* 2012;12:51.
 48. Vinsnes AG, Helbostad JL, Nyrønning S, Harkless GE, Granbo R, Seim A. Effect of physical training on urinary incontinence: a randomized parallel group trial in nursing homes. *Clin Intervent Aging* 2012;7:45-50.
 49. Wijk H, Corazzini K, Kjellberg IL, Kinnander A, Alexiou E, Swedberg K. Person-centered incontinence care in residential care facilities for older adults with cognitive decline: feasibility and preliminary effects on quality of life and quality of care. *J Gerontol Nurs* 2018;44(11):10-19.

50. Arnold EP, Milne DJ, English S. Conservative treatment for incontinence in women in rest home care in Christchurch: outcomes and cost. *Neurol Urodyn* 2015;35(5):636–641.
51. Ehlman K, Wilson A, Dugger R, Eggleston B, Coudret N, Mathis S. Nursing home staff members' attitudes and knowledge about urinary incontinence: the impact of technology and training. *Urolog Nurs* 2012;32(4):205–213.
52. Mathis S, Ehlman K, Dugger BR, Harrawood A, Kraft CM. Bladder Buzz: the effect of a 6-week evidence-based staff education program on knowledge and attitudes regarding urinary incontinence in a nursing home. *J Contin Ed Nurs* 2013;44(11):498–506.
53. Rahman AN, Schnelle JF, Applebaum R, Lindabury K, Simmons S. Distance coursework and coaching to improve nursing home incontinence care: lessons learned. *J Am Geriatr Soc* 2012;60(6):1157–1164.
54. Rahman AN, Schnelle JF, Osterweil D. Implementing toileting trials in nursing homes: evaluation of a dissemination strategy. *Geriatr Nurs* 2014;35(4):283–289.
55. Suzuki M, Iguchi Y, Igawa Y, Yoshida M, Sanada H, Miyazaki H, Homma Y. Ultrasound-assisted prompted voiding for management of urinary incontinence of nursing home residents: efficacy and feasibility. *Int J Urol* 2016;23(9):786–790.
56. Yu P, Hailey D, Fleming R, Traynor V. An exploration of the effects of introducing a telemonitoring system for continence assessment in a nursing home. *J Clin Nurs* 2014;23(21–22):3069–3076.
57. Moga DC, Carnahan RM, Lund BC, Pendergast JF, Wallace RB, Torner JC, Li Y, Chrischilles EA. Risks and benefits of bladder antimuscarinics among elderly residents of Veterans Affairs community living centers. *J Am Med Dir Assoc* 2013;14(10):749–760.
58. McClurg D, Cheater FM, Eustice S, Burke J, Jamieson K, Hagen S. A multi-professional UK wide survey of undergraduate continence education. *Neurourol Urodyn* 2013;32(3):224–229.
59. Morishita L, Uman GC, Pierson CA. Education on adult urinary incontinence in nursing school curricula: can it be done in two hours? *Nurs Outlook* 1994;42(3):123–129.
60. Australian Commission on Safety and Quality in Healthcare. Patient-centred care – improving quality and safety through partnerships with patients and consumers. Sydney: Australian Commission on Safety and Quality in Healthcare; 2011. Available from: <https://www.safetyandquality.gov.au/publications-and-resources/resource-library/patient-centred-care-improving-quality-and-safety-through-partnerships-patients-and-consumers>
61. Care Quality Commission. Regulation 9: person-centred care. UK: Care Quality Commission; 2022. Available from: <https://www.cqc.org.uk/guidance-providers/regulations-enforcement/regulation-9-person-centred-care>
62. Institute of Medicine. Crossing the quality chasm: a new health system for the 21st century. Washington (DC): National Academy Press (US); 2001.
63. Temkin-Greener H, Cai S, Zheng NT, Zhao H, Mukamel DB. Nursing home work environment and the risk of pressure ulcers and incontinence. *Health Serv Res* 2012;47:1179–200.
64. Yoon JY, Lee YJ, Bowers BJ, Zimmerman DR. The impact of organizational factors on the urinary incontinence care quality in long-term care hospitals: a longitudinal correlational study. *Int J Nurs Stud* 2012;49(12):1544–51.

Improving ward-based neurogenic bladder and bowel care after acute spinal cord injury: a narrative review

For referencing Levy C et al. Improving ward-based neurogenic bladder and bowel care after acute spinal cord injury: a narrative review. Australian and New Zealand Continence Journal 2023; 29(4):101-108.

DOI <https://doi.org/10.33235/anzcj.29.4.101-108>

Licensed under CC BY 4.0

Submitted 15 September 2023, Accepted 2 November 2023

ABSTRACT

Objectives There is a risk that early neurogenic bladder and bowel dysfunction following acute spinal cord injury (SCI) may not be adequately managed on the acute hospital wards. The aim of this paper was to review optimal assessment and management of patients with neurogenic bladder and/or bowel dysfunction during acute care immediately following SCI. A secondary aim was to translate these findings into an accessible policy for clinical staff involved in the care of acute SCI patients.

Methods A literature review was performed to identify the protocols and research describing the management of bladder and bowel dysfunction after acute SCI. The measures extracted were principles of practice, threshold measures and evaluation tools, and timelines of best care. Key findings relating to assessment, management and identification of the benefits of adhering to a protocol of clinical practice were summarised. Fidelity with evidence identified was evaluated, and practice gaps were recognised.

Results A total of 12 papers and six guidelines for providing excellent care to patients with SCI were identified. Overall, a systematic and comprehensive assessment of bladder and bowel function and neurological impairment should be completed early following SCI. Management is best individualised to both prevent known sequelae and optimise current function and quality of life. Acute SCI bladder and bowel management pathways were developed based on evidence identified.

Conclusions The study has identified that timely assessment and management of patients with spinal cord neurogenic bladder and/or bowel dysfunction during the acute phase is pivotal to optimising continence and mitigating risk of preventable harm prior to discharge to specialised spinal rehabilitation.

Keywords acute spinal cord injury, neurogenic bladder, neurogenic bowel, early spinal cord injury continence management pathway

BACKGROUND

Acute spinal cord injury (SCI) results from insults to the spinal cord and has permanent consequences, affecting independence and every aspect of life¹. Individuals may develop motor and sensory impairment and autonomic dysfunction caudal to the level of injury². The presence and severity of SCI is determined by the American Spinal Injury Association (ASIA) Impairment Scale (AIS) grade. In patients with an AIS grade A to D³ (which correlates with one or more impairments of light touch S4–5, pin prick S4–5, deep anal pressure S4–5 or voluntary anal contraction) elimination symptoms are common.

It should be noted that spinal shock lasts for around 3 months and is characterised by a loss of spinal reflexes and muscle tone below the level of injury⁴. Neurogenic bladder and/or bowel dysfunction during the period of spinal shock is not indicative of function in the chronic phase. The type of neurogenic bladder and/or bowel dysfunction is determined by the Neurological Level of Injury (NLI) and whether or not the sacral outflow is impacted.

Cristina Levy*

Department of Physiotherapy, The Royal Melbourne Hospital, VIC, Australia
Email cristina.levy@mh.org.au

Shilpa Gaikwad

Department of Neurosurgery, The Royal Melbourne Hospital, VIC, Australia

Michael Whishaw

Department of Aged Care and Sub-Acute Continence Service, The Royal Melbourne Hospital, VIC, Australia

Wendy Bower

Department of Physiotherapy, The Royal Melbourne Hospital, VIC, Australia

Department of Aged Care and Sub-Acute Continence Service, The Royal Melbourne Hospital, VIC, Australia
Faculty of Medicine, Dentistry and Health Science, University of Melbourne, VIC, Australia

*Corresponding author

Conflict of interest

The authors declare no conflicts of interest.

Funding

The authors received no funding for this study.

Neurogenic bladder dysfunction in patients with acute SCI is characterised by detrusor areflexia, impaired bladder sensation, non-relaxing bladder neck and proximal urethra, and possible impairment of pelvic floor muscle coordination⁴. Symptoms may include urinary frequency, urinary urgency, urinary incontinence⁴. Injuries below the conus medullaris (usually at T12 or below) usually lead to flaccid detrusor muscle and loss of innervation to the external urethral sphincter⁴. Filling of the bladder will occur at a low pressure; however, voiding is dysfunctional⁴. This results in bladder distension and high bladder volumes⁵. Injuries above the conus medullaris usually lead to spasticity of the muscles after resolution of spinal shock and are implicated in urinary tract infections and deterioration of the upper urinary tract⁵. Lesions at the level of the conus medullaris commonly cause pelvic floor dysfunction⁵.

Neurogenic bowel dysfunction occurs in 27–62% of patients with SCI and is characterised by dysfunction of the colon, abdominal muscles, pelvic floor muscles, rectum and external anal sphincter as well as impaired rectal and/or anal sensation^{6–8}. Symptoms may include constipation, abdominal distention and pain, bleeding (consequence of straining) and faecal incontinence⁶. Injuries below the conus medullaris generally lead to

flaccid pelvic floor muscles and external anal sphincter with no reflex response to increase intra-abdominal pressure. In addition, the tone of the external anal sphincter is affected⁷. The loss of descending inhibitory modulation significantly delays colonic transit time and may reduce compliance of the colon. Injuries above the conus medullaris lead to a reduced ability to effectively increase intra-abdominal pressure as well as an increased tone of the rectum and external anal sphincter⁸. Table 1 describes objective presentation and symptoms seen in the lower urinary tract and bowel after SCI.

The care for patients with SCI who have an acute neurogenic disruption to bladder and bowel function is complex and requires specialist knowledge and skills. Sequelae in both systems must be anticipated, recognised and managed appropriately in the acute phase. Prioritising this before patients arrive in post-acute rehabilitation minimises negative patterns and reduces the risk of complications developing^{4,8}.

The aim of this project was twofold. Firstly, to outline optimal assessment and management of patients with bladder and/or bowel dysfunction immediately following acute SCI based on research and best clinical practice. Secondly, to translate findings into a policy applicable to all clinical staff involved in the care of

Table 1. Important variables for bladder and bowel function in patients with SCI

Reflexive – spastic NLI above the conus medullaris (usually at T12)	Areflexive – flaccid NLI below the conus medullaris (usually T12 or below)
Bladder presentation	
<ul style="list-style-type: none"> • Intact micturition reflex • Increased tone of the detrusor muscle and external urethral sphincter • Impaired bladder sensation • Impaired voluntary control of the detrusor muscle and external urethral sphincter • High bladder pressures 	<ul style="list-style-type: none"> • Absent micturition reflex • Reduced tone of the bladder • Absent tone of the external urethral sphincter • Impaired bladder sensation • Impaired voluntary control of the detrusor muscle and external urethral sphincter • High bladder volume
Bladder symptoms	
<ul style="list-style-type: none"> • Urinary frequency • Urinary urgency • Urinary nocturia • Urine leakage • Bladder over-extension • Sexual dysfunction 	<ul style="list-style-type: none"> • Urinary overflow incontinence • Urinary retention • Bladder over-extension • Sexual dysfunction • Pelvic pain
Bowel presentation	
<ul style="list-style-type: none"> • Positive defecation reflex • Inability to effectively increase intra-abdominal pressure • Increased tone of the rectum and external anal sphincter • Impaired rectal sensation • Impaired voluntary control of the external anal sphincter 	<ul style="list-style-type: none"> • Absent defecation reflex • No reflex response to increase intra-abdominal pressure • Reduced tone of the rectum • Absent tone of the external anal sphincter • Impaired anal and rectal sensation • Impaired voluntary control of the external anal sphincter and pelvic floor muscles
Bowel symptoms	
<ul style="list-style-type: none"> • Faecal impaction proximal colon • Overflow incontinence • Haemorrhoids • (Less commonly) faecal incontinence 	<ul style="list-style-type: none"> • Faecal incontinence • Faecal impaction in the rectum • Overflow incontinence • Haemorrhoids

patients with acute spinal cord neurogenic bladder and/or bowel dysfunction.

METHODS

A literature review was performed for papers from the last decade that described management of bladder and bowel dysfunction after acute SCI. The databases of CINAHL Plus, Embase and MEDLINE were searched. Searches were restricted to English language only. A search strategy was developed that focussed on the following keywords as search terms: 'acute spinal cord injury', 'bladder or urinary bladder', 'bowel', 'incontinence or urinary incontinence or bowel incontinence', 'complications', 'assessment' and 'management'. Search terms were established by scoping using free text terms and exploding Medical Subject Headings (MeSH).

Studies in English and from the last decade that describe bladder and or bowel care early after any type of SCI were eligible for inclusion. No limits were placed on study types. Studies describing children or adolescents, neurological conditions aside from SCI or samples of SCI patients more than 3 months post-injury were excluded from the search. Information extracted were principles of practice, threshold measures and evaluation tools and timelines of best care.

All studies identified through the search results as describing assessment and treatment of bladder and bowel dysfunction after acute SCI were reviewed. Duplicate references were identified and removed. The titles and abstracts of the search results were screened, and full texts were retrieved for the relevant articles. Two reviewers completed this process independently. Any disagreements were resolved by consensus.

Articles were scrutinised for key findings related to systems, assessment, management and benefits of adhering to a protocol. Two authors independently identified and extracted data. No risk of bias assessment was performed as all study types were included. Current ward practice at our institution was benchmarked against variables identified. Findings were presented to the multidisciplinary team caring for patients with SCI and the evidence-practice gap between published regimes and routine assessment and management was discussed.

An optimal clinical pathway informed by study findings was formulated to align with the ward culture of our institution. This was then devolved into a policy that could be integrated into the local standards of care. The new policy was reviewed by the following key institutional stakeholders: director of neurosurgery, head of neurology, senior neurosciences pharmacist, urology clinical nurse consultants (CNCs), nursing unit managers for neurology and neurosurgery wards, policy coordinator and nurse educators. Feedback was integrated into the final documents and separate pathways for bladder and bowel care in the acute period following SCI were described.

RESULTS

In total, 31 papers were identified from the literature search; all were screened by two people. Following this, 12 papers⁹⁻²⁰ and six guidelines from SCI services²¹⁻²⁶ were relevant to the study aim and were included in the narrative review. The overall key findings were that a systematic and comprehensive assessment of bladder and bowel function and neurological impairment should be completed early following SCI before developing a management plan^{9,10,12,18,23}. Table 1 summarises the status of clinically important variables that should be evaluated in both systems to confirm the impact of SCI on bladder and bowel function.

It can be seen that the level of neurological injury dictates function and bladder and bowel symptoms. AIS grade and NLI (using the International Standards for Neurological Classification of SCI (ISNCSCI)) should be established to differentiate between three possible types of spinal cord neurogenic bowel and/or bladder dysfunction – reflexive (spastic), areflexive (flaccid) or mixed (spastic and flaccid) presentation. This should be re-assessed periodically as neurological improvement during admission may change the AIS grade. The Bulbocavernosus Reflex Test, reflecting the integrity of spinal reflex arcs at sacral levels 2-4, should be conducted serially to monitor for resolution of spinal shock. Additionally, a comprehensive post-SCI assessment should identify relevant medical details and medication use (particularly those that induce constipation), onset of autonomic dysreflexia and paralytic ileus. Evaluation will also establish fluid intake, diet and cognitive and functional capacity, for example hand skills and positioning.

Key findings related to management following acute SCI prioritised early education with patients and their family/carers to engage them in establishing an optimal management plan⁷. This education includes alteration in anatomy and physiology, understanding the need for ongoing assessment, management options, prognosis and rehabilitation planning¹⁴. Management is ideally individualised to both prevent known sequelae and optimise current function and quality of life.

With respect to bladder function, preservation of both the upper and lower urinary tracts are inter-related. Over-distention of the bladder, a trigger for autonomic dysreflexia, is prevented by insertion of an indwelling catheter (IDC). Transitioning from an IDC to intermittent catheterisation (IC) when the patient is medically stable is important to reduce the incidence of complications such as urinary traction infections (UTI)^{13,14}. Long-term suprapubic catheters (SPC) may be more suitable for patients with limited hand function or restricted carer support. Adherence to a daily routine minimises the risk of incontinence and UTIs. Multidisciplinary consultation will precede trial of void (TOV), ideally following the steps summarised in Figure 1¹⁷. Identification of detrusor pressures during bladder filling and smooth and musculoskeletal activity during filling and any emptying are clarified by urodynamic studies performed at least 6 weeks post-SCI or after resolution of spinal shock¹⁹.

Bowel management should similarly identify precursors of preventable changes (for example onset of paralytic ileus) and individualise intervention to avoid incontinence, optimise stool consistency and toileting at a suitable time and place, prevent complications, and enhance quality of life^{10,12}. This will include monitoring for signs of loose or hard stool, faecal impaction, faecal overflow incontinence, faecal incontinence and hyperkalaemia/hyponatremia. Digital rectal examination (DRE) should be performed to identify rectal fullness prior to inserting rectal stimulants or enemas and to remove rectally impacted faeces (DRF)¹¹. Follow-up DREs may be indicated at times but could potentially be replaced by small volume rectal washouts.

Six papers^{9,10,13-15,17} described the benefits of adhering to a protocol of clinical practice with respect to bladder and bowel function after acute SCI. An evidenced framework will clarify status, identify complications, prevent adverse changes (for example bladder over-distension), standardise transition of care components (e.g., from IDC to IC / the TOV process) and determine the need for specialist referrals for spinal rehabilitation. From a consumer perspective, educational content to support understanding of prognosis and informed decision-making based on the individual needs were highlighted.

Acute SCI bladder and bowel management pathways were developed to provide clear guidance to assist

1. Educate the patient to drink regularly, void as needed and to notify the nurses after each spontaneous micturition.
2. Perform a bladder scan 4–6 hours post-removal of the IDC.
3. Record three consecutive PVR volumes for up to 48 hours.
 - If the PVR is <200ml for three consecutive bladder scans and the patient is continent of urine, the bladder function is considered intact.
 - If the PVR is >200ml and/or the patient is incontinent of urine, the bladder is considered dysfunctional and an IC must be immediately completed or an IDC immediately re-inserted to empty the bladder completely.
4. If the patient does not void for 6 hours, use bladder scan to assess the urine volume.
 - If the bladder volume is >300ml, the bladder is considered dysfunctional and immediately re-insert an IC or an IDC to empty the bladder completely.
 - If the bladder volume is <300ml, the patient should be monitored for an additional 4 hours. If there is spontaneous micturition, continue the TOV process from step 3. If there is no spontaneous micturition, the bladder is considered dysfunctional and an IC or an IDC is re-inserted to empty the bladder completely.

If the patient who has been diagnosed with a dysfunctional bladder improves neurologically, the TOV process should be commenced.

IDC: indwelling catheter; PVR: post-void residual; IC: intermittent catheterisation; TOV: trial of void

Figure 1. Trial of void process for bladder management

clinicians with timely and appropriate assessment, management and treatment of patients with spinal cord neurogenic bladder and/or bowel dysfunction. The protocol was intended to be implemented within 24 hours of acute SCI and had utility for use by all clinicians involved in the care of patients with a SCI, eg, nurses, allied health clinicians, neurosurgical, neurology and urology medical clinicians and pharmacists. Separate pathways developed for bladder and bowel management are presented in Figures 2&3.

DISCUSSION

Functional limitations related to neurogenic bladder and bowel symptoms are profound, being ranked higher than loss of walking¹⁷. Not only can appropriate surveillance of bladder and bowel symptoms optimise continence, it can also identify adverse changes and prevent longer term complications. As described by the recently published *Australian and New Zealand clinical practice guidelines for the physiotherapy management of people with spinal cord injury*²⁷, the care of patients with SCI is complex, and bladder and bowel function is an important part of early rehabilitation. Bladder and bowel care in the presence of neurogenic disorders requires specialist knowledge and multidisciplinary skills. In the acute hospital setting, once SCI is diagnosed or suspected, neurogenic bladder and bowel dysfunction is anticipated and appropriate management commenced.

Our institution is not a recognised specialist service for people with SCI; however, spinal operations are frequently performed, generating a cohort of patients with some form of SCI. Non-specialist nurses, medical and allied health clinicians assess and plan treatment for SCI patients. As wait time for transfer to specialist spinal rehabilitation services is lengthy, patients with SCI often remain on the acute ward for an extended period. The risk of bladder or bowel complications developing during the patient’s acute admission should be appreciated. Benchmarking against evidence synthesised from the current literature review indicated that present practice on the acute wards in our non-specialised service does not align with best evidence. To address current gaps in practice recommendations, evidence gathered was tabulated as separate bladder and bowel management pathways.

Barriers and facilitators to the translation of these pathways into clinical processes of care within our institution’s unique culture and environment can be anticipated. Guihan¹⁵ reported the majority of barriers centred around current hospital workflow, policies, resources, education and training. Specifically, it is anticipated junior medical staff will require upskilling in performing ISNCSCI and Bulbocavernosus Reflex Tests and that nursing staff need more education on how to initiate and perform IC, particularly for males. This was identified by Goodes¹³ who noted that despite evidence supporting early implementation of IC, there were substantial delays in transitioning from IDC to IC due to staff training. Additionally, it is anticipated ward staff may be reluctant to complete invasive interventions




Figure 2. Bladder management in acute SCI pathway

AIS: ASIA (American Spinal Injury Association) Impairment Scale; NLI: Neurological Level of Injury; ISNCSCI: International Standards for Neurological Classification of SCI; IDC: indwelling catheter; TOV: trial of void; CNC: clinical nurse consultant; IC: intermittent catheterisation; SPC: suprapubic catheter




Figure 3. Bowel management in acute SCI pathway

AIS: ASIA (American Spinal Injury Association) Impairment Scale; NLI: Neurological Level of Injury; ISNCSCI: International Standards for Neurological Classification of SCI; IDC: indwelling catheter; DRE: digital rectal examination; DRF: digital removal of faeces

such as DRE and DRF. The development of educational in-services, training, online learning resources and targeted competencies will be implemented to address these barriers. The upskilling of clinicians is consistent with previous research demonstrating that in-service training for staff working in the acute SCI setting resulted in improved clinician knowledge, enhanced patient knowledge and job satisfaction¹⁰.

The introduction of the newly developed clinical pathways after acute SCI will fail unless the roll out follows implementation science principles. Prior to a change in practice, clinical staff will require access to education about the underlying rationale for modification and the opportunity to be part of an iterative process that optimises acceptability^{9,14}. The fit of the pathway process to the organisational culture should be negotiated, particularly whether additional tasks can be completed with current resources and technology. The documentation of the impact on patient status, for example the prevalence of complications after implementation, will ideally be embedded in electronic medical recording to facilitate audit and evaluation. In the interim, clinicians can endeavour to include more optimal continence strategies into their usual care of SCI patients. Future work will include the evaluation of educational components and co-design of optimal resources.

The strengths of this study include the benchmarking of current care against the latest evidence and clinical guidelines to provide a focus for change. Best practice management can potentially be individualised to target specific underlying dysfunction in each patient with acute SCI. The limitations of this work relate to our target as a single centre of care with unique culture and attributes and a low volume of acute SCI patients. The pathways require clinical champions to optimise implementation and a community of practice that prioritises bladder and bowel care in the acute phase of SCI.

In summary, this work has proactively addressed the issue of preventable complications in bladder and bowel function after acute SCI in a non-specialised hospital environment. Strategies that can be applied in the clinical context during patient assessment and management have been identified and integrated into system-specific multidisciplinary care pathways. Formal implementation to enhance continence care within our institution after SCI has been initiated, beginning with stakeholder consultation.

REFERENCES

- Lohne V, Severinsson E. Hope during the first months after acute spinal cord injury. *J Adv Nurs* 2004;47(3):279–286.
- Furlan JC, Fehlings MG. Cardiovascular complications after acute spinal cord injury: pathophysiology, diagnosis, and management. *Neurosurg Focus* 2008;25(5):E13.
- Alexander M, Carr C, Alexander J, Chen Y, McLain A. Assessing the ability of the Sacral Autonomic Standards to document bladder and bowel function based upon the Asia Impairment Scale. *Spinal Cord Ser Cases* 2019;5:85.
- Cardozo L, Rovner E, Wagg A, Wein A, Abrams P, editors. *Incontinence*, 7th edition. ICI-ICS International Continence Society; 2023. Chapter 4, A. Initial assessment of urinary incontinence in adult male and female patients B. Patient-Reported Outcome Assessment, pp.308–9.
- Taweel WA, Seyam R. Neurogenic bladder in spinal cord injury patients. *Res Rep Urol* 2015 Jun 10;7:85–99.
- Beuret-Blanquart F, Weber J, Gouverneur JP, Demangeon S, Denis P. Large bowel transit time and anorectal manometric abnormalities in 19 patients with complete transection of the spinal cord. *J Auton Nerv Syst* 1990;25:109–112.
- Krogh K, Mosdal C, Gregersen H, Laurberg S. Rectal wall properties in patients with acute and chronic spinal cord lesions. *Dis Colon Rectum* 2002;45(5):641–649.
- Lynch AC, Antony A, Dobbs BR, Frizelle FA. Bowel dysfunction following spinal cord injury. *Spinal Cord* 2001;39(4):193–203.
- Bragge P, Guy S, Boulet M, Ghafoori E, Goodwin D, Wright B. A systematic review of the content and quality of clinical practice guidelines for management of the neurogenic bladder following spinal cord injury. *Spinal Cord* 2019;57(7):540–549.
- Borsh S, Sikka S, Callender L, Bennett M, Reynolds M, Driver S. Implementation of a neurogenic bowel program for spinal cord injury in the acute care setting: perceptions of patients and staff. *Occ Ther Health Care* 2019;33(3):306–322.
- Coggrave MJ, Norton C. The need for manual evacuation and oral laxatives in the management of neurogenic bowel dysfunction after spinal cord injury: a randomized controlled trial of a stepwise protocol. *Spinal Cord* 2010;48(6):504–510.
- Cotterill N, Madersbacher H, Wyndaele JJ, Apostolidis A, Drake MJ, Gajewski J, Heesakkers J, Panicker J, Radziszewski P, Sakakibara R, Sievert KD, Hamid R, Kessler TM, Emmanuel A. Neurogenic bowel dysfunction: clinical management recommendations of the Neurologic Incontinence Committee of the Fifth International Consultation on Incontinence. *Neurourol Urodyn* 2018;37(1):46–53.
- Goodes LM, King GK, Goodwin DM, Watts A, Bardley J, Middleton J, Bragge P, Dunlop SA. Barriers and facilitators to optimising inpatient bladder management after spinal cord injury. *Spinal Cord* 2020;58:1291–1300.
- Goodwin D, Brock J, Dunlop S, Goodes L, Middleton J, Nunn A, Wright B, Bragge P. Optimal bladder management following spinal cord injury: evidence, practice and a cooperative approach driving future directions in Australia. *Arch Phys Med Rehab* 2018;99(10):2118–2121.
- Guihan M, Bosshart HT, Nelson A. Lessons learned in implementing SCI clinical practice guidelines. *SCI Nurs* 2004;21(3):136–142.
- Krassioukov A, Eng JJ, Claxton G, Sakakibara BM, Shum S. Neurogenic bowel management after spinal cord injury: a systematic review of the evidence. *Spinal Cord* 2010;48(10):718–733.
- Lim V, Mac-Thiong JM, Dionne A, Begin J, Richard-Denis A. Clinical protocol for identifying and managing bladder dysfunction during acute care after traumatic spinal cord injury. *J Neurotrauma* 2021;38(6):718–724.
- Sekido N, Igawa Y, Kakizaki H, Kitta T, Sengoku A, Takahashi S, et al. Clinical guidelines for the diagnosis and treatment of lower urinary tract dysfunction in patients with spinal cord injury. *Int J Urol* 2020;27(4):276–288.

19. Welk B, Schneider MP, Thavaseelan J, Traini LR, Curt A, Kessler TM. Early urological care of patients with spinal cord injury. *World J Urol* 2018;36(10):1537-1544.
20. Yeung HY, Iyer P, Pryor J, Nicholson M. Dietary management of neurogenic bowel in adults with spinal cord injury: an integrative review of literature. *Disabil Rehab* 2021;43(9):1208-1219.
21. Birrer K, Hobbs B, Giancarelli A, Ashworth S, Clancy R. Acute spinal cord injury management. Orlando Regional Medical Center; 2018 [cited 2022 Nov 4]. Available from: <http://www.surgicalcriticalcare.net/Guidelines/Acute%20Spinal%20Cord%20Injury%202018.pdf>
22. Byrne C, Clements R, Mahony A. Victorian Spinal Cord Service - neurogenic bladder management. Austin Health; 2018.
23. Coggrave MJ. Multidisciplinary Association of Spinal Cord Injured Professionals. Guidelines for management of neurogenic bowel dysfunction in individuals with central neurological conditions. MASCIP Middlesex; 2012 [cited 2022 Nov 4]. Available from: <https://www.mascip.co.uk/wp-content/uploads/2015/02/CV653N-Neurogenic-Guidelines-Sept-2012.pdf>
24. Hogan M, Mullane J. Management of spinal neurogenic bladder & bowel. University College London Hospitals NHS Foundation Trust; 2019.
25. Pryor J, Fisher M, Middleton J. Management of the neurogenic bowel for adults with spinal cord injuries. NSW Agency for Clinical Innovation; 2014 [cited 2022 Nov 4]. Available from: https://aci.health.nsw.gov.au/__data/assets/pdf_file/0019/155215/Management-Neurogenic-Bowel.pdf.
26. Spiteri R, Brown A. Victorian Spinal Cord Service - neurogenic bowel management. Austin Health; 2009.
27. Glinsky JV, Harvey LA, and the Australian and New Zealand Physiotherapy Clinical Practice Guidelines consortium. Australian and New Zealand Clinical Practice Guideline for the physiotherapy management of people with spinal cord injury. 2022 [cited 2023 May 7]. Available from: <https://sciptguide.com/wp-content/uploads/2023/07/Australian-and-NZ-SCI-Physiotherapy-Clinical-Practice-Guidelines-1.1-1.pdf>

Abstracts

UroGynaecological Society of Australasia Annual Scientific Meeting Melbourne 22-25 November 2023

We are pleased to publish a small selection of abstracts with permission from the authors. For the full abstract program contact UGSA www.ugsa.com.au

Clinical Quality Indicators using Australasian Pelvic Floor Procedure Registry clinical data

Aruna Kartik

Australasian Pelvic Floor Procedure Registry (APFPR), Melbourne, Australia.

DOI <https://doi.org/10.33235/anzcj.29.4.109>

Licensed under CC BY 4.0

Introduction The Australasian Pelvic Floor Procedure Registry (APFPR) was established to monitor the safety and quality of Stress Urinary Incontinence (SUI) and Pelvic Organ Prolapse (POP) procedures involving mesh or other pelvic floor prostheses.

Aims The APFPR has developed initial Clinical Quality Indicators (CQIs) to benchmark, inform and support improvements in quality of care. This presentation reports results from APFPR's first iteration of CQIs derived from its dataset. These CQIs aim to demonstrate APFPR's potential for supporting quality improvement.

Methods Potential CQIs were derived from process and outcome measures based on the Australian Commission on Safety and Quality in Health Care guidance, and the APFPR dataset. Inclusion and exclusion criteria were developed in conjunction with the APFPR Clinical Advisory Committee. A descriptive analysis of APFPR procedure data was undertaken. CQIs were reported for SUI, and POP with combined POP+SUI cohorts, and will be presented overall and as funnel plots.

Results As of 08 August 2023, the APFPR comprises approximately 500 patient procedures, with 23 participating sites reporting the following CQIs.

The proportion of initial patient procedures (SUI/POP) with:

- objective clinical assessments completed (90.2/91.8%)
- intraoperative cystoscopy performed (99.2/100%)
- improved patient outcomes (82.6/95.2%)
- the proportion of initial/subsequent patient procedures for SUI and POP with:
 - a return to theatre (0.3/0%)
 - readmission within 30 days (3.5/2%)
 - catheterisation on discharge (5.2/0%).

Discussion The development of CQIs is an important milestone for the APFPR. Initial CQIs have been designed to capture common processes and outcomes. Further CQIs that incorporate patient-reported outcomes will be developed in 2024.

Source of Funding Commonwealth Department of Health and Aged Care.

Biomechanical and immunobiological properties of human fascia lata (HFL) vs TiMesh®: implications for pelvic reconstructive surgery

David Hennes

Monash University, Melbourne, Australia.

DOI <https://doi.org/10.33235/anzcj.29.4.110>

Licensed under CC BY 4.0

Introduction The use of autologous human fascia lata (HFL) in pelvic reconstructive procedures such as sacrocolpopexy (SCP) has become increasingly desired by patients, due to a greater awareness of potential complications of mesh and restrictions on their use in Urogynaecology.

Aims This study aimed to compare the morphometric and biomechanical properties, and in vivo foreign body response of HFL with TiMesh®, in a pre-clinical murine abdominal incision model.

Methods HFL was harvested from women undergoing sacrocolpopexy or pubovaginal sling insertion (n=26). HFL and TiMesh® were characterised using uniaxial tensiometry. HFL or TiMesh® was implanted in C57/BL6 immunocompetent mice via an abdominal skin incision (n=8 mice/gp/time-point). Explanted tissue analysis at 7 and 90 days included histology and electron microscopy for cellular infiltration, angiogenesis, elastin and collagen content. Quantitative fluidigm PCR assessed 80 genes associated with ECM homeostasis, cell adhesion, angiogenesis and inflammation. Data analysis was conducted using the Mann-Whitney test in GraphPad Prism version 7.0.

Results HFL consists of acellular fibrous collagen, exhibiting superior tissue integration and host cell infiltration to TiMesh®, which demonstrated loose integration and increased acute inflammation. HFL had higher resistance to passive repetitive loading and absolute tensile strength in comparison with TiMesh®. At 90 days, there was a significant increase in genes associated with ECM production (Tgfrb1), ECM regulation (Mmp2) and angiogenesis (Fgf1, Ang-1) in the HFL group compared with TiMesh®, indicating significantly improved wound healing.

Discussion HFL is an ideal alternative surgical graft for augmentative urogynaecological procedures, with superior mechanical adaptability, durability and tissue integration compared to TiMesh®.

Source of Funding Cabrini Research Foundation Grant (\$30,000).

Mesh perforation into viscus following pelvic mesh surgery: Experiences and recommendations for diagnosis and management

Yu Hwee Tan

Queensland Pelvic Mesh Service, Gold Coast Health and Queensland Pelvic Floor Services, Greenslopes Private Hospital, Queensland, Australia.

DOI <https://doi.org/10.33235/anzcj.29.4.111>

Licensed under CC BY 4.0

Introduction Pelvic mesh has been used for the treatment of pelvic organ prolapse (POP) and stress urinary incontinence (SUI). Between 3-20% of women suffer from complications with debilitating long-term outcomes.

Aims Provide recommendations regarding diagnosis and management of pelvic mesh perforation into viscus.

Methods A retrospective observational study of patients who had diagnosis and management of mesh in viscus at three tertiary urogynaecology units was performed.

Results Fifty-eight patients were diagnosed with mesh in viscus following cystoscopy and examination under anaesthesia (EUA). Mesh involved included mid-urethral slings - retropubic (36.9%), transobturator (18.5%) and single-incision slings (10.8%), transvaginal POP mesh (15.4%), abdominal sacrocolpopexy (13.8%), and uncertain type (4.6%). Viscus involved included bladder (39.7%), urethra (50%), bladder and urethra (3.4%), and rectum (6.9%). Presenting symptoms included mixed urinary incontinence [75.9%], recurrent UTIs (48.3%), voiding dysfunction (19%), and pain (56.9%). Fifty patients underwent mesh excision and viscus repair (with or without concomitant labial fat flap), with successful repair in all patients (100%). 48% had complete mesh excision. Mean follow up was 9.5 months (range 0.5 - 96 months). Post-operatively, thirty-three patients (63.5%) had recurrent urinary incontinence, seven (13.5%) had persistent pain and seven (13.5%) had recurrent UTIs. None had lower genitourinary tract fistula or wound breakdown.

Discussion All patients required EUA and cystourethroscopy for diagnosis. Excision of mesh in viscus appears to reduce risk of recurrent mesh erosion or fistula. Consideration should be given to performing a labial fat flap during urethral repair to prevent injury from future treatment for persistent pelvic floor dysfunction.

NEWS

AUSTRALIAN NEWS

In the final edition of the *ANZCJ* for 2023, I am proud to report on all we have achieved at the Continence Foundation of Australia. Overall, 2023 has been a busy year for the Foundation as we continue to advocate and advance the voice of continence in Australia. The year has noted several substantial projects, including multi-channel campaigns for World Continence Week and the Great Dunny Hunt, professional development events including the National Conference on Incontinence, and successful collaborations with Jean Hailes and the Prostate Cancer Foundation of Australia. The past quarter has been busy for the Foundation, with numerous events and partnerships coming to fruition.

Women's Health Week: 4–8 September 2023

In support of Women's Health Week, the Foundation has continued our community partnership with Jean Hailes to inspire women of all ages to focus on good health and wellbeing.

This year, Jean Hailes had five daily themes under the banner 'grow your knowledge', encouraging women to make informed decisions about their health with topics including pain drain, hello hormones and food for thought. Aligning with these themes, the Foundation focused on reminding women of all ages there is much they can do to reduce their risk of incontinence. In fact, in many cases, incontinence may be prevented, better managed or even cured.

A highlight of the week was an encore screening of the Lived Experience panel webinar, where five panellists shared their open and honest experiences with managing incontinence, barriers to seeking help and the common misconceptions they have faced. Facilitated by award-winning disability advocate Anja Christoffersen, around 100 attendees registered to watch the webinar. The Foundation also launched episode four of the new podcast series *This is my story* which featured guest Stephanie Thompson and a discussion regarding childbirth injury and pelvic organ prolapse. Please visit continence.org.au/womens-health-week-resources to access these resources.

State education days

During September and October the State and Territory Advisory Committees (STACs) organised education days. These education days posed an informative professional development, networking and learning opportunity for health professionals across Australia and overseas working in bladder, bowel and pelvic floor health. Program highlights included a paediatric pelvic health workshop held at the Perth Children's Hospital in Western Australia and a series of interactive workshops in New South Wales which covered practical bowel management, continence surgery and urodynamics. With close to 100 attendees at each event and nearly 60 exhibitors, this year's state education days were

informative and inspiring. According to a delegate at the Victoria education day:

As always this was an amazing event. Such a wealth of knowledge shared by like-minded professionals. I really enjoyed the presentations and having an opportunity to meet others, network and just have a chat. Thank you so much for a wonderful day. I feel motivated and refreshed. Looking forward to the next event.

If you would like to hear about other professional development events, sign up to our [events mailing list](#).

Physiotherapy special interest group events

The Physiotherapy Special Interest Group (SIG) was introduced as a national initiative in 2023. With over 110 members, the physiotherapy SIG hosts specialist education events and offers strong network opportunities for practitioners with varying levels of clinical experience from across Australia.

In October the SIG hosted a webinar titled *Exercise during pregnancy and return to sport* with a focus on sharing the latest research updates in the fields of exercise during pregnancy, return to sport and diastasis management. The webinar attracted nearly 200 registrants from across Australia and overseas. Visit cfaphysios.com.au if you would like to learn more about the SIG and becoming a member.

BINS4Blokes collaboration with the Prostate Cancer Foundation of Australia

BINS4Blokes, a men's health initiative of the Continence Foundation of Australia, has recently teamed up with the Prostate Cancer Foundation of Australia (PCFA) to help get more continence product disposal bins in male public toilets across Australia. "We're proud to partner with BINS4Blokes to help raise awareness of the need for continence product disposal bins and encourage more councils, businesses and shopping centres to include these bins in male public toilets", said Bernard Riley, PCFA head of supportive care programs.

Over 25,000 men are diagnosed with prostate cancer every year and many of these people will experience temporary or long-term continence issues after treatment. The goal of the partnership is to create a strong advocacy network of those with lived experience who can support the installation of incontinence product disposal bins in male toilets in their local community.

This collaboration has already seen positive return, with a prostate cancer survivor and PCFA support group member successfully advocating for the installation of incontinence product disposal bins in the Ballina Shire Council.

Education

Our education team is always busy keeping healthcare professionals up to date with the latest information on

bladder and bowel continence, with online learning modules and free webinars.

Currently the Foundation has 37 different education courses and individual modules, as well as other professional development opportunities. This is inclusive of both paid and free courses as well as free supported learning modules (such as Continence Resources for Aged Care).

The recordings from the joint 31st National Conference on Incontinence and the 4th Functional Urology Symposium Conference 2023 have also now been uploaded onto the eLearning platform, allowing all registrants to review presentations from this year's event.

I am looking forward to 2024 as we continue our vision and purpose of an Australian community free from the restrictions of incontinence across the lifespan.

Rowan Cockerell
CEO, Continence Foundation of Australia

31st National Conference on Incontinence & 4th Functional Urology Symposium



Thank you to our sponsors and partners

Supported by







Principal partners

Gold sponsors





Silver sponsors







Bronze sponsors

NEW ZEALAND NEWS

Continence NZ Conference 2023

After a hiatus of almost a decade, it was our pleasure to host our much-anticipated National Conference in Auckland in September 2023. We welcomed almost 100 people working in continence care for 2 days to discuss the latest research and trends in continence care for both adults and children. We are so grateful for the support of our event sponsors, Auckland Unlimited, Lion Foundation, SPQ Limited and Confitex, as well as the speakers and exhibitors who featured at the event. After such a long period with no face-to-face events or education due to COVID-19, it was fantastic to reconnect with our members and industry partners. Visit continencenzconference.org for photos from the event; we thank you for your support.

Community and online education

Our popular Toilet Tactic webinar series ran from August to November 2023 with children's continence nurse Lisa Smith. The series covers topics such as bowel and bladder health, toilet training, constipation, bedwetting, daytime wetting, stool withholding and toilet training for children with additional needs. It's free to register for one or all our sessions. We will have further webinars for both adults and children running throughout the year. Visit continence.org.nz/pages/Upcoming-Community-Education/301 for details of our webinars.

Online children's continence clinics for the Auckland region

In response to a large number of calls to the Continence NZ helpline, Continence NZ is piloting a new offering of online clinics. These clinics are for whānau (families) living in the Auckland area with children experiencing issues with toilet training, constipation, stool withholding, and daytime or night-time wetting. Families receive limited support for these issues through their local GP and services such as Plunket. These are the first clinics of their kind for Continence NZ and consist of an information session for six to eight whānau, then a one-on-one follow up with a children's continence nurse.

Pregnancy guide

Continence NZ has developed a guide to pelvic health before and during pregnancy which will be launched over the summer. Pregnancy and childbirth are an ideal opportunity for risk assessment and prevention strategies to be implemented with women because research indicates that this life stage corresponds with major risk factors for pelvic floor dysfunction. New Zealand does not currently have enough pelvic health physiotherapists working in publicly funded healthcare to support every pregnant woman. The costs associated with private pelvic health physiotherapy are out of reach of many expectant whānau. The guide will provide comprehensive, clear and practical information about bladder, bowel and pelvic health before,

during and after pregnancy for all birthing parents. Visit continencenz.org.nz for further information.

Research update

Last summer, Continence NZ engaged with nearly 300 members of our community in the most comprehensive research project we have undertaken to date. The research aim was to better understand the experiences of those living with incontinence in Aotearoa and how we can support them. We engaged with our members, health professionals, people living with incontinence and their whānau to understand their experiences and how Continence NZ can help.

The research highlighted incontinence as a growing issue in New Zealand, which impacts the quality of life of nearly 1.25 million New Zealanders. The environmental cost of incontinence product disposal is estimated at NZ\$14.9 million per year. Incontinence can be improved or resolved in many cases with the aid of continence nurse specialists and pelvic health physiotherapists. However, many New Zealanders are not receiving the treatment they need due to shortages in this specialist care. An estimated investment of NZ\$6.4 million per year into pelvic health nurses and pelvic health physiotherapists would result in NZ\$8.9 million in savings, producing a net benefit of NZ\$2.5 million per year. Importantly, it would enable New Zealanders with incontinence to continue to live their best lives. This is achieved by lessening the economic, social and personal impact that is often associated with incontinence. We are incredibly grateful to all those who participated in the research. Visit communityresearch.org.nz/research/continence-nz-service-users-research-and-service-review to see the full research.

Laura Fear
CEO, Continence NZ